



Thermoregulation in long distance race: Literature review

Termorregulação em corridas de longa distância: Revisão bibliográfica

DOI: 10.56238/isevjhv2n6-010

Receipt of originals: 01/12/2023

Publication Acceptance: 12/18/2023

Igor Gomes Breguez Rocha

Professor of the Discipline of Physiology at the University of the UNC Contestado
Campus Porto União Santa Catarina
E-mail: igorbreguez@gmail.com

Marco Antonio Schueda

Coordinator of the Faculty of Medicine and Professor of the Discipline of Anatomy at the
University of Contestado UNC Campus Porto União - Santa Catarina
E-mail: schueda.sc@gmail.com

Anna Júlia Pelchebiski

Undergraduate student of the Medicine course at the University of Contestado UNC
Campus Porto União - Santa Catarina
E-mail: annajuliapelchebiski@gmail.com

Beatriz Ribeiro da Silva de Amorim

Undergraduate student of the Medicine Course at the University of Contestado UNC Campus
Porto União - Santa Catarina
Email: beatriz.medicina2023@gmail.com

Rafaella Gugelmin Barros

Undergraduate student of the Medicine Course at the University of
Contested UNC Campus Porto União - Santa Catarina
E-mail: rafaella.gugelmin@gmail.com

Thaysa Bruna Gavronski

Undergraduate student of the Medicine Course at the University of
Contested UNC Campus Porto União - Santa Catarina
E-mail: thaysa101@hotmail.com

Study carried out at the Discipline of Physiology, Faculty of Medicine, Universidade do
Contestado - UNC, Porto União - Santa Catarina

ABSTRACT

The literature review of the present study demonstrates that during physical exercise, the body produces heat through the active muscles, generating heat stress. Through the analysis of articles mainly from the last five years, we conclude that to keep body temperature stable, the body uses regulation mechanisms dependent on environmental, task, and personal parameters. In intense exercise and in hot environments, blood flow to the skin can increase significantly, and sweating is also elevated. However, non-thermal reflexes during exercise cause the regulation of blood flow in the skin to be different from that at rest. The body adjusts the threshold for the dilation of the skin's blood vessels to a higher core body temperature and modifies the sensitivity to this



temperature change. Regular training improves these adaptations, allowing athletes to dissipate heat more effectively. This review highlights the physiological mechanisms involved in thermal regulation during exercise, discusses the estimation of core temperature in humans, and presents new and nutritional approaches to assess parameters to keep core temperature stable.

Keywords: Physiology, Body temperature, Heat stress.

1 INTRODUCTION

The body's core temperature is normally set to 36.6°C ¹, but it can diverge considerably when exposed to extreme conditions. The lowest recorded core body temperature with survival by a human is 13.7°C ². In contrast, during exercise in the heat, well-trained athletes can reach core body temperatures of 41.5°C ³. Core body temperature in humans is the main variable regulated in thermoregulation⁴. Resting core body temperature also depends on age, gender, ethnicity, ambient temperature, humidity, time of day and month of the year¹. The amplitude of this diurnal variation is 0.5°C in healthy individuals⁵, excluding any exposure to extreme cold or heat stress, fever or exercise. The elderly have a lower resting core temperature than young adults^{6,7,8}. The menstrual cycle significantly alters core body temperature, with an upward change of 0.4°C during the luteal phase compared to the follicular phase in premenopausal women^{9,10}.

The skin acts as an interface with the environment, but unlike core temperature, skin temperature is not regulated¹¹ and varies throughout the body in response to the thermal environment^{12,13}. Even so, the average skin temperature can be categorized as cold ($<30^{\circ}\text{C}$), hot ($30\text{--}34.9^{\circ}\text{C}$) and hot ($>35^{\circ}\text{C}$)¹⁴. Data indicate that changes in core temperature are much smaller than those in the skin and extremities in relation to changes in ambient temperature^{15,16}.

Thermoregulation mechanisms include sweating, peripheral vasodilation, tachycardia, and increased respiration¹⁶.

Sweating is responsible for dissipating heat through the evaporation of sweat on the surface of the skin.

Peripheral vasodilation increases blood flow to the skin, allowing heat to dissipate through radiation and conduction.

Tachycardia and increased breathing help increase metabolic rate and heat dissipation. In addition, the hypothalamus is the main control center for thermoregulation in the human body, monitoring body temperature and activating these mechanisms to maintain thermal homeostasis.

We will explore these mechanisms in the literature review below.



2 METHODOLOGY

The methodology chosen to carry out the following research was a literature review with an exploratory approach.

For Marconi and Lakatos (2019), literature reviews have the purpose of putting the researcher in direct contact with everything that has been written, said or filmed on the subject. It is not a mere repetition of the subject, but provides analysis of the theme from another point of view or approach, in order to reach new conclusions¹⁷.

The research followed the recommended methods, after choosing the theme and preliminary research. The samples, were read, selected, evaluated and analyzed. The characteristics of the research were defined and then the results were discussed, interpreted and presented.

The guiding question was: How does body temperature regulation occur during physical activity in long-distance running?

For a bibliographic review of the theme, searches were carried out in printed bibliographies and digitized articles. The articles were collected, preferably, in the last five years, surveying studies specifically related to the proposed theme, which resulted in 83 articles/chapters.

In the researched articles, a pattern was noticed in the problem addressed, that is, the authors also had the same doubt as in the present study with the object of the research.

Understanding this subject is of great importance for medical professionals as well as physical educators and physical therapists.

3 RESULTS AND DISCUSSION

3.1 AUTONOMIC AND BEHAVIORAL THERMOREGULATION

The regulation of body temperature is carried out through parallel processes of behavioral and autonomic thermoregulation.

Behavioral regulation of temperature operates largely through conscious behavioral adjustments, which when in the heat include a range of coolness-seeking behaviors, such as staying in the shade, drinking cold drinks, pouring water on the head, and wearing light-colored clothing. It has also been suggested that adjustments in the rate of work during exercise in the heat constitute behavioral adjustments that contribute to regulating body temperature¹⁸.

Thermoregulatory behavior decreases the need for autonomic responses¹⁹, which operate through physiological processes independent of conscious voluntary behavior. These responses include control of vasomotor (i.e., cutaneous vasodilation) and sudomotor (sweating) function in



heat, along with metabolic heat production (i.e., shivering) and vasomotor function (i.e., cutaneous vasoconstriction) in cold. Environments. A negative feedback system is typically described as the regulatory system that mediates thermoregulatory autonomic responses. This physiological control system produces graduated responses according to disturbances in a regulated variable: core body temperature. Such control structures are called proportional control systems. In humans, central (brain, spine, and gastrointestinal tract) and peripheral (skin) thermoreceptors provide afferent input to thermoregulatory centers located in the hypothalamus, where they are compared with the set point^{20,21}. The set point is purely a mathematical concept used to describe the thermal control of effector responses and does not imply a specific neural model of thermoregulation or defined temperature. Instead, it describes different stages of recruitment within the magnitude of a load error, which is the difference between the input and the set point²². Thus, central and peripheral thermoreceptors send information to a central integrator, located in the preoptic anterior hypothalamus²³. The notion of central integration is supported by data describing the proportion of contributions of core temperature and skin inputs to sudomotor control^{24,25} and vasomotor^{24,26,27}. These responses can also be altered by factors such as circadian rhythm, fever, phase of the menstrual cycle, and acclimatization to heat^{22,28}.

The notion of central integration has evolved, with the emergence of other regulatory models. These include a model in which thermal equilibrium is achieved by regulating heat across a range of heat loads, detecting heat flow to and from the body, and defending body heat content through thermoeffector responses^{29,30}. Another model suggests that independent thermoeffector circuits coordinate their activities to regulate body temperature around an equilibrium point^{11,31}. Thermoregulation during exercise in the heat is regulated in a similar way to that at rest and influenced by factors such as hydration status and environmental conditions, as well as work rate (i.e., exercise intensity)¹⁶.

3.2 THERMAL EQUILIBRIUM

Human heat equilibrium refers to the balance between the internal rate of metabolic heat production and the rate of heat exchange with the surrounding environment through sensitive (i.e., convection, conduction, and radiation) and insensitive (i.e., evaporation) pathways.

Four main environmental parameters affect the biophysical properties of human thermal equilibrium: ambient temperature, humidity, air velocity, and solar radiation. In addition to environmental factors, task-dependent parameters have an impact on heat exchange: metabolic

heat production rate and clothing; as well as personal parameters: body surface area, body mass, gender, age and aerobic fitness¹⁶.

3.2.1 Environmental parameters

Ambient temperature, humidity, air velocity, solar radiation

3.2.2 Task-Dependent Parameters

Metabolic heat production, clothing

3.2.3 Personal parameters


Body surface area, body mass, sex, age, and aerobic fitness

3.3 ENVIRONMENTAL PARAMETERS:

3.3.1 Ambient temperature

During exercise, ambient temperatures higher than skin temperature led to sensible (i.e., dry) heat gain, while lower temperatures led to heat loss (FIGURE 1). Environments with high ambient temperature and low humidity favor heat loss through evaporation, since sweat and mucosal moisture can evaporate more easily¹⁶.

Fig. 1 - As the ambient temperature increases and approaches the skin temperature, dry heat loss is reduced and evaporation becomes the main route of heat dissipation¹⁶.



3.3.2 Humidity

Absolute humidity refers to the amount of water vapor present in the air. In the atmosphere, absolute humidity ranges from almost zero to 30g/m³ when the air is saturated at 30°C. Air humidity is strongly related to climate. Hot, humid climates are typically found in rainforest areas,



and hot, dry climates near deserts. High absolute humidity compromises the evaporation capacity of sweat from the skin because the difference in water vapor (i.e., humidity) between the skin surface and the environment is low¹⁶.

3.3.3 Air Velocity

The standard meteorological wind speed is determined at 10 m above the ground and the highest air speed ever recorded on Earth is 113.3 m/s. Air speed during exercise depends on factors such as the direction of travel, wind direction, and terrain. The displacement of air through the body results in convective heat exchange, depending on the thermal gradient between the air and the skin. Air displacement also helps with heat loss through evaporation, as it removes the layer of saturated water vapor that can stagnate in the skin¹⁶.

3.3.4 Solar radiation

The amount of solar radiation that reaches the human body depends on the location on Earth, the time of day, the season, and the level of the exposed skin area.

The integration of environmental parameters can be used to provide an index of the severity of the thermal environment, with a combination of parameters used to calculate different indices. A meta-analysis identified more than 300 thermal indices, of which 185 were included in various statistical analyses³². Temperature (98%), relative humidity (RH: 77%) and air velocity (72%) were the most commonly used environmental parameters to calculate the severity of the environment, with solar radiation (45%) and a combination of the four parameters (42%) being less frequently integrated into these calculations. The most commonly used index in sport and exercise is the moistbulb globe. temperature index (WBGT), which has been advocated by the American College of Sports Medicine³³ and the National Athletic Trainers' Association³⁴ and used by several international sports organizations (e.g., World Athletics, World Triathlon).

3.4 TASK-DEPENDENT PARAMETERS.

3.4.1 Metabolic heat production

Human metabolism is the sum of resting metabolism (65 W/m²) and exercise. Oxidation of substrates during exercise contributes significantly to increasing the body's core temperature, since only 20 to 25%^{35,36} The metabolic energy is converted into mechanical work, most of which is released as heat. The highest maximum oxygen uptake rate ever recorded (VO₂max) is 96.7 ml/kg/min³⁷. Although not sustainable over a very long period, this level of oxygen consumption



is equivalent to 2,500 W of metabolic heat production, which underlines the large contribution of exercise-induced heat production to human thermal balance¹⁶.

3.4.2 Garments (clothing)

Clothing acts as a barrier between the skin and the environment, altering heat exchange properties in relation to environmental conditions. The properties of the material and the fit of a garment can affect thermal stress during exercise, reducing heat dissipation and promoting heat conservation³⁸. As such, the insulating properties and water vapour resistance of garments worn while exercising in the heat should be as low as possible. The water absorption capacity of the material should also be low, as the sweat trapped in the clothing does not evaporate and does not provide cooling. In addition, the reflective properties of a garment are important in high radiative load scenarios (i.e., direct sunlight). It is not so much the color of the clothing that is important, but the reflective properties of the dyes used in the clothing. Ventilation in the air layer between the skin and clothing (i.e., bellows effect) is important for heat loss during exercise in the heat³⁹. An example of the bellows effect is seen in the desert, where no difference in skin temperature was observed when wearing loose black or white clothing (i.e., robes), despite a 60C difference in the surface temperature of clothing (black: 47 C; white: 41 C)^{40,41,42}.

3.5 PERSONAL PARAMETERS.

3.5.1 Body Surface Area

The heat generated during metabolism is lost on the surface of the body to prevent excessive heat storage. Body surface area is calculated using a formula based on height and weight: surface area = $0.20247 \text{ height (m)}^{0.725} \text{ weight (kg)}^{0.42543}$, which has been validated using three-dimensional scanning techniques⁴⁴. A large body surface area is beneficial for heat loss through evaporation, as the number of active sweat glands is proportional to the surface area⁴⁵. Dry heat loss is also increased by having a larger body surface area when the ambient temperature is lower than the skin temperature⁴⁶. Thus, for a given thermal environment, the potential for heat loss is greater in those with a large body surface area⁴⁷.

3.5.2 Body Surface Area Ratio

Individuals with a high body surface area-to-mass ratio experience less heat storage during exposure to noncompensable heat than those with a lower ratio, due to the larger area for dry and evaporative heat loss relative to body mass⁴⁷. The body surface/mass ratio decreases with



increasing body mass and to a greater extent in women⁴⁸. Thus, for heavy women it is more difficult to release body heat than for equally heavy men. This was confirmed in a study in which a thermal model was used to assess core body temperature in women of different body morphologies, showing that relatively fat women achieve considerably higher body temperatures during exercise in a hot/humid environment than their thinner women⁴⁹.

3.5.3 Sex

Men and women differ in body size, with men generally being heavier, taller, and exhibiting higher VO₂max. However, when standardized for body surface area, metabolic heat production during various tasks is similar between genders⁵⁰.

3.5.4 Age

Aging impacts both thermoregulatory capacity and fluid regulation^{51,52}. Older individuals (>60 years) have a lower resting core body temperature, attenuated cutaneous vasodilator capacity, less effective sweat response, and decreased thermoreceptor sensitivity compared to younger individuals⁵¹⁻⁵³. The elderly also have a higher thirst threshold^{54,55}, lower total body water level^{56,57}, reduced renal function^{56,58} and impaired regulation of plasma vasopressin at rest and after dehydration^{59,60}. These regulatory functions deteriorate with advancing age⁶¹ and increase the risk of developing hyperthermia and dehydration^{51,52,60,62,63}. However, older individuals in good shape retain better thermoregulation capacity and can improve thermoregulatory capacity with training⁵⁹.

3.5.5 Aerobic fitness

Regular resistance exercise leading to improved aerobic fitness (i.e., VO₂max) has been shown to increase the ability to lose heat. Aerobic training activates cutaneous vasodilation at a lower core temperature and increases skin blood flow to a certain core temperature⁶⁴⁻⁶⁶. The increase in blood flow to the skin is largely mediated by the expansion of blood volume and higher cardiac output that characterize the training state⁶⁷. Resistance training has also been reported to lower the internal temperature threshold for the onset of sweating, increase the sweat rate at a given core temperature, and increase the maximum sweat rate^{65,68}. Modelling suggests that an increase in VO₂max induced by exercise training of 12 to 17% should reduce the internal temperature threshold for sweat onset by 0.1°C⁶⁹. However, the improved sweating function associated with aerobic fitness may also be related to regular resistance training, providing a



repeated thermal challenge that leads to improvements in thermoregulatory capacity^{70,71}. Core body temperature can be measured at different locations on the body, depending on the type of equipment and need for accuracy.

3.5.6 Exertional heat illness (EHI)

Heat production often exceeds the heat dissipation capacity. Although the thermophysiological responses to exercise in the heat are well understood, individual responses vary substantially^{72,73}. The severity of HIE varies along its continuum, with mild complaints after exercise-associated muscle cramps, to more serious concerns during heat syncope and heat exhaustion, and life-threatening risks during heat stroke^{34,74-76}.

Dehydration and intake of specific medications (i.e., diuretics) may further increase the risk of heat syncope, while blood volume expansion induced by heat acclimatization may reduce the risk.

The fact that exertional heat stroke continues to be the third leading cause of death in athletes in the United States, after cardiac disorders and head and neck trauma^{77,78}, further emphasizes the need for early recognition of HIE symptoms by health professionals, appropriate equipment to treat heat stroke victims (i.e., ice baths) and education of race participants. Studies considering heart rate and sweat rate, two practical (non-invasive) variables, have shown greater predictive power in the prevention of HIE⁷⁹.

3.6 WATER BALANCE CORPORAL¹⁶

The total volume of body water represents approximately 60% of body mass (range: 45 to 75%) and depends on age and sex, with lower values for the elderly and women. Body water can be divided into intracellular and extracellular fluid compartments. The intracellular compartment accounts for approximately (~) 40% of body mass, while the extracellular compartment can be segmented into the interstitial (~15% of body mass) and intravascular (i.e., plasma volume, ~5% of body mass) subcompartments.

When moderate- to high-intensity exercise is initiated, a vasoconstrictor-mediated reduction in skin blood flow occurs, which contributes to providing additional blood to working skeletal muscles. Shortly thereafter, cutaneous vasodilation is initiated to aid in the dissipation of metabolically generated heat. Splanchnic and renal vasoconstriction may contribute 600 to 800 ml/min⁻¹ of blood to this response. The internal temperature threshold at which active cutaneous



vasodilation occurs is linked to exercise intensity, with an increase in work rate delaying the threshold relative to rest.

The change in cutaneous vasodilation threshold during exercise results in lower cutaneous blood flow for a given core temperature compared to rest. This response is exacerbated when exercising in cold environmental conditions compared to hot environmental conditions, as lower skin temperatures suppress the active vasodilator response to increased internal temperature, thereby increasing the temperature threshold for vasodilation. Factors such as time of day, phase of the menstrual cycle in women, and plasma osmolality have also been shown to influence the threshold of onset of cutaneous vasodilation.

Acute dehydration is associated with decreased glycemic regulation, worsening mood, blunting of blood pressure control, reduced cerebral blood flow during sympathoexcitation, and orthostatic intolerance.

Hypohydration and dehydration may reduce the responsiveness of the central neural structures that regulate heat loss by evaporation, with an increase in sweating onset threshold associated with a hyperosmolality-induced inhibition of heat-sensitive neurons within the hypothalamus (i.e., median preoptic nucleus). In analysis, body water losses not replenished (i.e., hypohydration) or inadequately replaced (i.e., dehydration) during exercise under heat stress reduce blood volume and increase osmolality, with the severity of these responses impacting thermoregulatory capacity.

3.7 NUTRITION^{80,81,82}

The primary source of energy for humans is carbohydrate (CHO), which is made up of molecules of carbon, hydrogen, and oxygen atoms. They are classified into: monosaccharides, oligosaccharides and polysaccharides, whose energy value is equal to that of proteins, which provide 4 kcal for every⁸⁰ grams.

Despite being the main source of energy, they have a variety of beneficial effects on the body, including those on hormone and neurotransmitter production, metabolism, metabolic efficiency, insulin sensitivity, and gut microbiota⁸⁰.

Carbohydrates are categorized based on the degree of structural complexity, are ingested in the form of complex molecules (starch) or simple molecules (glucose) and invariably transformed by digestion into monosaccharides (simple carbohydrates): glucose, fructose and galactose⁸⁰.



The importance of using carbohydrates as a source of energy during the practice of various sports has been recognized since the beginning of the last century. This macronutrient is the largest source of energy in the human diet, representing about 40 to 80% of the total energy consumed, and its use as an ergogenic resource to increase sports performance is indicated because it is a determining substrate in increasing performance⁸⁰.


The first known study demonstrating the usefulness of glucose as a key energy source for long-distance runners was conducted in 1924. Marathon runners developed hypoglycemia with exhaustion, according to the study. Since then, several studies have stated that glucose is an important macronutrient for maintaining performance in long-term activities⁸⁰.

Consuming carbohydrates during exercise is a common practice in many sports, especially endurance sports like running. When compared to a regular (50%) and low-carbohydrate (10%) diet, a high-carbohydrate diet (70% of CHO's dietary energy) and large muscle glycogen stores appears to increase endurance capacity. Although knowledge of human physiology and nutrition, as well as dietary changes and nutrient supplementation.

When carbohydrate-rich foods are consumed, they reach the stomach, where they are broken down and a product called glucose is released through a process called gluconeogenesis^{81,82}. In anaerobic metabolism, the body uses only glycogen for energy through the creation of glucose^{81,82}. This process, called glycolysis, produces energy quickly, but only two molecules of ATP (energy) are obtained^{81,82}. On the other hand, aerobic metabolism uses fat and glycogen as energy and creates 38 ATP molecules through a more time-consuming process^{81,82}. The aerobic and anaerobic systems occur in separate cellular compartments (mitochondria and cytoplasm, respectively) and often at different rates, involve different reactants and products^{81,82}. It is not only of interest to allocate the two separate energy processes, but also which fuel is being used^{81,82}. Glucose and fatty acids provide most of the fuel needed for energy production in skeletal muscles during aerobic exercise, while glucose is the main source of energy in anaerobic exercise.

The body has significantly more energy available in the form of fat, but the utilization rate of this form of energy cannot be increased at high intensities of exercise when anaerobic metabolism is the primary mechanism^{81,82}. Thus, the body is able to use mainly fatty acids as a source of energy at low levels of intensity^{81,82}. When not exercising, 30% of the body's energy comes from glycogen and 70% from fat reserves^{81,82}. These percentages change as the intensity increases, as does the number of calories burned. Glucose is preferred because it is readily available and rapidly metabolized, but it is limited.

FIG. 2 - Fuel Utilization: allocation of calories per minute per kilogram used as a function of percent VO₂max^{81,82}



It is also well established that an increase in exercise intensity mediates an increase in glycogenolysis and carbohydrate oxidation and a decrease in fat oxidation. This progressive change in the mobilization and utilization of the energy substrate during exercise in hot environmental conditions may represent the maintenance of a gradually increasing relative exercise intensity, mediated by a hyperthermia-induced decrease in maximal aerobic capacity¹⁶.

3.7.1 Contemporary nutritional strategies to optimize performance in long-distance runners and walkers⁸³

Distance events in athletics include cross country, 10,000 m track race, half marathon and road marathon and 20 km race and 50 km race walking events in different terrain and environmental conditions.

Race times for elite athletes range from 26 minutes to >4 hours, with the key factors for success being high aerobic power, the ability to exercise at a large fraction of that power, and high running/walking economy. Nutrition-related contributors include body mass and anthropometry, ability to use fuels, particularly carbohydrates (CHO) to produce adenosine triphosphate economically over the duration of the event, and maintenance of

reasonable hydration status in the face of sweat losses induced by exercise intensity and the environment. Race nutrition strategies include CHO-rich eating in the hours of the days leading up to the event to store glycogen in sufficient amounts for the event's fuel needs, and in some cases, CHO and fluid consumption during the race to make up for losses in events.



Beneficial CHO intake ranges from small amounts, including mouthwash in the case of shorter races, to high intake rates (75–90 g/h) in longer runs. A personalized and practiced nutrition plan should balance the benefits of the fluids and CHO consumed within the practical opportunities, in relation to the time, cost, and risk of intestinal discomfort. In hot environments, hyperhydration or pre-race cooling strategies can provide a small but useful compensation for accumulated thermal challenge and fluid deficit. Sports foods (beverages, gels, etc.) can help in training/race meetings nutrition plans, with caffeine and perhaps nitrate being used as evidence-based performance supplements⁸³.

3.7.2 Thermoregulatory Function Adjustments¹⁶

The rate of total body sweat during physical activity is determined by exercise intensity and weather conditions, but typically ranges between 0.5 and 2.0 L·h⁻¹, with some athletes (~2%) sweating >3.0 L·h⁻¹. Gradual reductions in body mass (e.g., 2 to 5%) due to fluid deficit result in marked reductions in plasma ($\geq 10\%$) and blood ($\geq 6\%$) volume. Loss of plasma volume with exercise leads to a state of hyperosmotic hypovolemia that is proportional to the decrease in total body water. Although the composition of the precursor secretory fluid contained in eccrine sweat glands is similar to that of plasma, a considerable amount of ions (e.g., sodium and chloride) within the fluid are reabsorbed as it passes through the gland duct, leading to a sweat osmolality of approximately half that of plasma. The increase in intravascular osmotic pressure resulting from plasma hyperosmolality causes a change in fluid from the intracellular to the extracellular compartment, which helps to defend the plasma volume. However, at high levels of total body sweat, the volume of fluid mobilized from the intracellular compartment to the vasculature is insufficient to restore plasma volume and leads to intracellular dehydration¹⁶.

Increased core temperature during exercise in the heat with hypohydration (2 to 7% loss of body mass). Several studies have shown that hyperosmotic hypovolemia induced by hypohydration delays thermoregulatory sweating and cutaneous vasodilation at rest and during exercise and reduces the sensitivity of the relationship between thermoeffector responses and core temperature. Although both hypovolemia and hyperosmolality influence these responses, the increase in plasma osmolality seems to be more strongly correlated with the reduction of sweating during exercise-heat stress than the decrease in blood volume^{A.17}.

4 CONCLUSION



To keep body temperature stable, the body uses regulation mechanisms dependent on environmental, task, and personal parameters. In intense exercise and in hot environments, blood flow to the skin can increase significantly, and sweating is also elevated. However, non-thermal reflexes during exercise cause the regulation of blood flow in the skin to be different from that at rest. The body adjusts the threshold for the dilation of the skin's blood vessels to a higher core body temperature and modifies the sensitivity to this temperature change. Regular training improves these adaptations, allowing athletes to dissipate heat more effectively.



REFERENCES

- Obermeyer Z, Samra JK, Mullainathan S. Individual differences in normal body temperature: longitudinal big data analysis of patient records. *BMJ*359: 359:j5468, 2017. doi:10.1136/bmj.j5468.
- Gilbert M, Busund R, Skagseth A, Nilsen PÅ, Solbø JP. Resuscitation from accidental hypothermia of 13.7C with circulatory arrest. *Lancet*355: 375–376, 2000. doi:10.1016/S0140-6736(00)01021-7.
- Racinais S, Moussay S, Nichols D, Travers G, Belfekih T, Schumacher YO, Periard JD. Core temperature up to 41.5C during the UCI Road Cycling World Championships in the heat. *Br J Sports Med*53: 426–429, 2019. doi:10.1136/bjsports-2018-099881.
- Benzinger TH. Heat regulation: homeostasis of central temperature in man. *Physiol Rev*49: 671–759, 1969. doi:10.1152/physrev.1969.49.4.671.
- Krauchi K, Wirz-Justice A. Circadian rhythm of heat production, heart rate, and skin and core temperature under unmasking conditions in men. *Am J Physiol Regul Integr Comp Physiol*267: R819–829, 1994. doi:10.1152/ajpregu.1994.267.3.R819.
- Fox RH, Woodward PM, Exton-Smith AN, Green MF, Donnison DVMH. Body temperatures in the elderly: a national study of physiological, social, and environmental conditions. *Br Med J*1:200–206, 1973. doi:10.1136/bmj.1.5847.200.
- Gomolin IH, Aung MM, Wolf-Klein G, Auerbach C. Older is colder: temperature range and variation in older people. *J Am Geriatr Soc*53: 2170–2172, 2005. doi:10.1111/j.1532-5415.2005.00500.x.
- Kenney WL. Thermoregulation at rest and during exercise in healthy older adults. *Exerc Sport Sci Rev*25: 41–76, 1997.
- Baker FC, Waner JI, Vieira EF, Taylor SR, Driver HS, Mitchell D. Sleep and 24 hour body temperatures: a comparison in youngmen, naturally cycling women and women taking hormonal contra-ceptives. *J Physiol*530: 565–574, 2001. doi:10.1111/j.1469-7793.2001.0565k.x.
- Cagnacci A, Arangino S, Tuveri F, Paoletti AM, Volpe A. Regulation of the 24h body temperature rhythm of women in luteal phase: role of gonadal steroids and prostaglandins. *Chronobiol Int*19: 721–730, 2002. doi:10.1081/CBI-120005394.
- Romanovsky AA. Thermoregulation: some concepts have changed. Functional architecture of the thermoregulatory system. *Am J Physiol Regul Integr Comp Physiol*292: R37–R46, 2007. doi:10.1152/ajpregu.00668.2006.
- Bierman W. The temperature of the skin. *JAMA*106: 1158–1162, 1936. doi:10.1001/jama.1936.02770140020007.
- Gagge AP, Gonzalez RR. Mechanisms of heat exchange: biophysics and physiology. In: *Handbook of Physiology: Environmental Physiology*. Bethesda, MD: American Physiological Society, 1996, p. 45–84.
- Sawka MN, Cheuvront SN, Kenefick RW. High skin temperature and hypohydration impair aerobic performance. *Exp Physiol*97:327–332, 2012. doi:10.1113/expphysiol.2011.061002.
- Olesen BW. Thermal comfort. *Bruel and Kjaer Technical Review*2:3–41, 1982.
- Périard JD, Eijssvogels TMH, Daanen HAM. Exercise under heat stress: thermoregulation, hydration, performance implications, and mitigation strategies. *Physiol rev.* vol 101, october 2021, pg 1873-1979



Marconi MA, Lakatos EM. Fundamentos de Metodologia Científica. São Paulo, SP, 8ª ed. Atlas, pg. 200

Flouris AD, Schlader ZJ. Human behavioral thermoregulation during exercise in the heat. *Scand J Med Sci Sports*25: 52–64, 2015. doi:10.1111/sms.12349.

Schlader ZJ, Vargas NT. Regulation of body temperature by auto-nomic and behavioral thermoeffectors. *Exerc Sport Sc Rev*47: 116–126, 2019. doi:10.1249/JES.000000000000180.

Hammel HT, Jackson DC, Stolwijk JA, Hardy JD, Stromme SB. Temperature regulation by hypothalamic proportional control with an adjustable set point. *JApplPhysiol*18: 1146–1154, 1963. doi:10.1152/jappl.1963.18.6.1146.

Hensel H. Neural processes in thermoregulation. *Physiol Rev*53:984–1017, 1973. doi:10.1152/physrev.1973.53.4.948.

Gisolfi CV, Wenger CB. Temperature regulation during exercise: old concepts, new ideas. *Exerc Sport Sci Rev*12: 339–372, 1984.

Boulant JA. Hypothalamic neurons regulating body temperature. In: *Handbook of Physiology: Environmental Physiology*, edited by Fregly MJ, Blatteis CM. New York: Oxford Press, 1996, p.105–125.

Bleichert A, Behling K, Scarperi M, Scarperi S. Thermoregulatory behavior of man during rest and exercise. *Pflugers Arch*338: 303–312, 1973. doi:10.1007/BF00586072.

Wyss CR, Brengelmann GL, Johnson JM, Rowell LB, Niederberger M. Control of skin blood flow, sweating, and heart rate: role of skin vs. core temperature. *J Appl Physiol*36: 726–733, 1974. doi:10.1152/jappl.1974.36.6.726.

Olofsson P. Increasing mean skin temperature linearly reduces the core-temperature thresholds for vasoconstriction and shivering in humans. *Anesthesiology*82: 1160–1168, 1995. doi:10.1097/0000542-199505000-00011.

Frank SM, Raja SN, Bulcao CF, Goldstein DS. Relative contribution of core and cutaneous temperatures to thermal comfort and auto-nomic responses in humans. *J Appl Physiol*86: 1588–1593, 1999. doi:10.1152/jappl.1999.86.5.1588.

Stephenson LA, Kolka MA. Effect of gender, circadian period and sleep loss on thermal responses during exercise. *Human Performance Physiology and Environmental Medicine at Terrestrial Extremes*, edited by Pandolf KB, Sawka MN, Gonzalez RR. Indianapolis, IN: Benchmark Press, 1988, p.267–304.

Bazett HC. Theory of reflex controls to explain regulation of body temperature at rest and during exercise. *J Appl Physiol*4: 245–262, 1951. doi:10.1152/jappl.1951.4.4.245

30 - Webb P. The physiology of heat regulation. *Am J Phys Regul Integr Comp Physiol*268: R838–R850, 1995. doi:10.1152/ajpregu.1995.268.4.R838.

Kobayashi S. Temperature-sensitive neurons in the hypothalamus: a new hypothesis that they act as thermostats, not as transducers. *Prog Neurobiol*32: 103–135, 1989. doi:10.1016/0301-0082(89)90012-9.

Loannou LG. Thermal indices and occupational heat stress: a systematic review and meta-analysis. In: *Effects of Heat on Behavioral and Physiological Mechanisms of the Human Thermoregulatory System during Rest, Exercise, and Work (PhD Thesis)*. Thessaly, Greece: University of Greece, chapt 6, 2020.



Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, Pyne SW, Roberts WO. American College of Sports Medicine position stand. Exertional heat illness during training and competition. *Med Sci Sports Exerc*39: 556–572, 2007. doi:10.1249/MSS.0b013e31802fa199.

Casa DJ, DeMartini JK, Bergeron MF, Csillan D, Eichner ER, Lopez RM, Ferrara MS, Miller KC, O'Connor F, Sawka MN, Yeargin SW. National Athletic Trainers' Association position statement: exertional heat illnesses. *J Athl Train*50: 986–1000, 2015. doi:10.4085/1062-6050-50.9.07.

Kristoffersen M, Sandbakk O, Ronnestad BR, Gundersen H. Comparison of short-sprint and heavy strength training on cycling performance. *Front Physiol*10: 1132, 2019. doi:10.3389/fpls.2019.01132, 10.3389/fphys.2019.01132.

Lucía A, Hoyos J, Perez M, Santalla A, Chicharro JL. Inverse relationship between VO₂max and economy/efficiency in world-class cyclists. *Med Sci Sports Exerc*34: 2079–2084, 2002. doi:10.1249/01.MSS.0000039306.92778.DF

Ronnestad BR, Hansen J, Stenslokken L, Joyner MJ, Lundby C. Case Studies in Physiology: Temporal changes in determinants of aerobic performance in individual going from alpine skier to world junior champion time trial cyclist. *J Appl Physiol*127: 306–311, 2019. doi:10.1152/jappphysiol.00798.2018.

Daanen HA. Physiological strain and comfort in sports clothing. In: *Textiles for Sportswear*, edited by Shishoo R. Amsterdam, The Netherlands: Elsevier/Woodhead Publishing, 2015, p.153–168.

Havenith G. Interaction of clothing and thermoregulation. *Exog Dermatol*11: 221–230, 2002. doi:10.1159/000068802.

Shkolnik A, Taylor CR, Finch V, Borut A. Why do Bedouins wear black robes in hot deserts? *Nature*283: 373–375, 1980. doi:10.1038/283373a0.

Moran-Cortes, J.F.; Gómez-Martín, B.; Escamilla-Martínez, E.; Sánchez-Rodríguez, R.; Gómez-Carrión, Á.; Martínez-Nova, A. Thermoregulation in Two Models of Trail Run Socks with Different Fabric Separation. *Life* 2023, 13, 1768. <https://doi.org/10.3390/life13081768>

Di Domenico I, Hoffmann SM, Collins PK. O papel das roupas esportivas na termorregulação, conforto e desempenho durante exercícios no calor: uma revisão narrativa. *Sports Med - Aberto* 8, 58 (2022). <https://doi.org/10.1186/s40798-022-00449-4>

DuBois D, DuBois EF. Clinical calorimetry: Tenth paper a formula to estimate the approximate surface area if height and weight be known. *Arch Intern Med (Chic)*XVII: 863–871, 1916. doi:10.1001/archinte.1916.00080130010002.

Tikuissis P, Meunier P, Jubenville CE. Human body surface area measurement and prediction using three-dimensional body scans. *Eur J Appl Physiol*85: 264–271, 2001. doi:10.1007/s004210100484.

Lee JB, Park TH, Lee HJ, Yun B. Sex-related differences in sudomotor function in healthy early twenties focused on activated sweat gland density. *Chin J Physiol*63: 1, 2020. doi:10.4103/CJP.CJP_46_19.

Notley SR, Park J, Tagami K, Ohnishi N, Taylor NA. Morphological dependency of cutaneous blood flow and sweating during compensable heat stress when heat-loss requirements are matched across participants. *J Appl Physiol*121: 25–35, 2016. doi:10.1152/jappphysiol.00151.2016.

Cramer MN, Jay O. Biophysical aspects of human thermoregulation during heat stress. *Auton Neurosci*196: 3–13, 2016. doi:10.1016/j.autneu.2016.03.001.



Robinette KM, Daanen H, Paquet E. The CAESAR project: A 3-D sur-face anthropometry survey. In: Proceedings–2nd International Conference on 3-D Digital Imaging and Modeling, 3DIM 1999. Washington, DC: IEEEComputer Society, 1999, p. 380–386.

Yokota M, Berglund LG, Bathalon GP. Female anthropometric variability and their effects on predicted thermoregulatory responses to work in the heat. *Int J Biometeorol*56: 379–385, 2012. doi:10.1007/s00484-011-0441-6.

Zhai Y, Li M, Gao S, Yang L, Zhang H, Arens E, Gao Y. Indirect calorimetry on the metabolic rate of sitting, standing and walking officeactivities. *Build Environ*145: 77–84, 2018. doi:10.1016/j.buildenv.2018.09.011.

Blatteis CM. Age-dependent changes in temperature regulation–amini review. *Gerontol*58: 289–295, 2012. doi:10.1159/000333148.

Chester JG, Rudolph JL. Vital signs in older patients: age-related changes. *J Am Med Dir Assoc*12: 337–343, 2011. doi:10.1016/j.jamda.2010.04.009.

Guergova S, Dufour A. Thermal sensitivity in the elderly: a review. *Ageing Res Rev*10: 80–92, 2011. doi:10.1016/j.arr.2010.04.009.

Mack GW, Weseman CA, Langhans GW, Scherzer H, Gillen CM, Nadel ER. Body fluid balance in dehydrated healthy older men: thirst and renal osmoregulation. *J Appl Physiol*76:1615–1623,1994. doi:10.1152/jappl.1994.76.4.1615.

Phillips PA, Rolls BJ, Ledingham JG, Forsling ML, Morton JJ, Crowe MJ, Wollner L. Reduced thirst after water deprivation in healthy elderly men. *N Engl J Med*311: 753–759, 1984. doi:10.1056/NEJM198409203111202.

Davis KK. Disorders of fluid balance: dehydration and hyponatremia. In: *Principles of Geriatric Medicine and Gerontology*, edited by Hazard WB, Blass JP, Ettinger WH, Halter JB. New York: McGrawHill, 1994, p.1182–1190.

Schoeller DA. Changes in total body water with age. *Am J ClinNutr*50: 1176–1181, 1989. doi:10.1093/ajcn/50.5.1176.

Lindeman RD, Tobin J, Shock NW. Longitudinal studies on the rate of decline in renal function with age. *J Am Geriatr Soc*33: 278–285, 1985. doi:10.1111/j.1532-5415.1985.tb07117.x.

Miller JH, Shock NW. Age differences in the renal tubular responseto antidiuretic hormone. *JGerontol*8: 446–450, 1953. doi:10.1093/geronj/8.4.446.

Sheehy CM, Perry PA, Cromwell SL. Dehydration: biological considerations, age-related changes, and risk factors in older adults. *BiolRes Nurs*1: 30–37, 1999. doi:10.1177/109980049900100105.

Bongers CC, Eijsvogels TM, Nyakayiru J, Veltmeijer MT, ThijssenDH, Hopman MT. Thermoregulation and fluid balance during a 30-km march in 60- versus 80-year-old subjects. *Age (Dordr)*36:9725, 2014. doi:10.1007/s11357-014-9725-1.

Begum MC.A review of the literature on dehydration in the institu-tionalized elderly.*Eur J Clin*5: 47–53, 2010.

Mentes J.Oral hydration in older adults: greater awareness isneeded in preventing, recognizing, and treating dehydration. *Am JNurs*106: 40–49, 2006. doi:10.1097/00000446-200606000-00023.



Beaudin AE, Clegg ME, Walsh ML, White MD. Adaptation of exercise ventilation during an actively-induced hyperthermia following passive heat acclimation. *Am J Physiol Regul Integr Comp Physiol*297: R605–R614, 2009. doi:10.1152/ajpregu.90672.2008.

Roberts MF, Wenger CB, Stolwijk JA, Nadel ER. Skin blood flow and sweating changes following exercise training and heat acclimation. *J Appl Physiol Respir Environ Exerc Physiol*43: 133–137, 1977. doi:10.1152/jappl.1977.43.1.133.

Thomas CM, Pierzga JM, Kenney WL. Aerobic training and cutaneous vasodilation in young and older men. *J Appl Physiol* (1985)86:1676–1686, 1999. doi:10.1152/jappl.1999.86.5.1676.

Simmons GH, Wong BJ, Holowatz LA, Kenney WL. Changes in the control of skin blood flow with exercise training: where do cutaneous vascular adaptations fit in? *Exp Physiol*96: 822–828, 2011. doi:10.1113/expphysiol.2010.056176.

Henane R, Flandrois R, Charbonnier JP. Increase in sweating sensitivity by endurance conditioning in man. *J Appl Physiol Respir Environ Exerc Physiol*43: 822–828, 1977. doi:10.1152/jappl.1977.43.5.822.

Havenith G. Individualized model of human thermoregulation for the simulation of heat stress response. *J Appl Physiol* (1985)90:1943–1954, 2001. doi:10.1152/jappl.2001.90.5.1943.

Ravanelli N, Gagnon D, Imbeault P, Jay O. A retrospective analysis to determine if exercise training-induced thermoregulatory adaptations are mediated by increased fitness or heat acclimation. *Exp Physiol*106: 282–289, 2021. doi:10.1113/EP088385.

Calegari A, Souza RM. Termorregulação e desempenho esportivo em condições de calor extremo. *evinci, unibrasil*, volume 9, número 2, p. 1e2, outubro, 2023.

Veltmeijer MT, Thijssen DH, Hopman MT, Eijsvogels TM. Within subject variation of thermoregulatory responses during repeated exercise bouts. *Int J Sports Med*36: 631–635, 2015. doi:10.1055/s-0034-1398676.

Westwood CS, Fallowfield JL, Delves SK, Nunns M, Ogden HB, Layden JD. Individual risk factors associated with exertional heat illness: a systematic review. *Exp Physiol*106: 191–199, 2021. doi:10.1113/EP088458.

Leon LR, Bouchama A. Heat stroke. *Compr Physiol*5: 611–647, 2015. doi:10.1002/cphy.c140017.

Sawka MN, Leon LR, Montain SJ, Sonna LA. Integrated physiological mechanisms of exercise performance, adaptation, and maladaptation to heat stress. *Compr Physiol*1: 1883–1928, 2011. doi:10.1002/cphy.c100082.

Sonna LA, Sawka MN, Lilly CM. Exertional heat illness and human gene expression. *Prog Brain Res*162: 321–346, 2007. doi:10.1016/S0079-6123(06)62016-5.

Howe AS, Boden BP. Heat-related illness in athletes. *Am J Sports Med*35: 1384–1395, 2007. doi:10.1177/0363546507305013.

Barrow MW, Clark KA. Heat-related illnesses. *Am Fam Physician*58: 749–756, 759, 1998.

Andrade MT, Nunes-Leite MMS, Bruzzi RS, Souza CH, Uendele-Pinto JP, Prado LS, Soares DD, Gonçalves DAP, Coimbra CC, Wanner SP. Predicting the body core temperature of recreational athletes at the end of a 10 km self-paced run under environmental heat stress. *Experimental Physiology*, 108, 852–864 2023. <https://doi.org/10.1113/EP091017>



Santos,CN, Denadai LKV, Cruz MFA. The importance of carbohydrate for sports performance in running athletes: an integrative literature review. *Research, Society and Development*, v. 12, n. 6, e19712642260, 2023

(CC BY 4.0) | ISSN 2525-3409 | DOI: <http://dx.doi.org/10.33448/rsd-v12i6.42260>

Cook C, Chen G, Hager WW, Lenhart S. Optimally controlling nutrition and propulsion force in a long-distance running race. *Front. Nutr.* 10:1096194 2023. doi: 10.3389/fnut.2023.1096194

Stipanuk MH, Caudill MA. *Biochemical, Physiological and Molecular Aspects of Human Nutrition*. 3rd ed. St. Louis, MO: Elsevier/Saunders (2013).

Burke LM, Jeukendrup AE, Jones AM, Mooses M. Contemporary Nutrition Strategies to Optimize Performance in Distance Runners and Race Walkers. *Int J Sport Nutr Exerc Metab.* 2019 Mar 1;29(2):117-129. doi: 10.1123/ijsnem.2019-0004. Epub 2019 Apr 4. PMID: 30747558.