



Microbiological analysis of salads consumed raw in the cafeteria of an educational institution

10.56238/isevmjv3n4-001

Receipt of originals: 06/12/2024

Acceptance for publication: 07/02/2024

Wéltima Teixeira Cunha¹.

INTRODUCTION

The National School Feeding Program (PNAE) establishes criteria for the management of school meals, the School Feeding Council, canteens and kitchens in schools, and the work of nutritionists and educators at school (CONCEIÇÃO, 2015; BRASIL, 2009a).

In this sense, the PNAE, implemented in 1955, is a public policy and aims to transfer financial resources to ensure school meals for students in all basic education, early childhood education, high school and youth and adult education enrolled in public and philanthropic schools, to meet the nutritional needs of these students during their stay in the classroom, contributing to growth, the development, learning and school performance of students, and focuses on the formation of healthy eating habits (BRASIL, 2009b; CECANE PARANÁ, 2010).

The consumption of vegetables is essential and should be part of any family's menu. It occurs due to its nutritional value arising from vitamins, minerals, fibers, low caloric intake and the fibers contained in them (NASCIMENTO et al., 2005). It is noteworthy that the general population has changed its eating habits with regard to the consumption of fresh vegetables (OLIVEIRA et al., 2006).

It should be noted, however, that vegetables consumed raw are responsible for the transmission of enteric diseases. Contamination occurs in the vegetable garden, by the use of irrigation water or inadequate fertilizers, in transport or by handling at points of sale; and the successive manipulations increase the chances of contamination (TAKAYANAGUI, 2001).

This contamination is due to the presence of a group of coliforms and one of them is bacteria from the total coliform group. The other group of fecal coliforms (coliforms at 45° C or thermotolerant) in food indicates the hygienic-sanitary conditions of the product and a better indication of the possible presence of enteropathogens (FRANCO; LANDGRAF, 2008).

The concern with the quality and safety of food is a global public health issue, due to the

¹ Federal Institute of Education, Science and Technology of Bahia - Bahia
weltimateixeira@ifba.edu.br



possibility of ingesting some type of food contaminated by pathogenic microorganisms (CUNHA, 2006). Vegetables that are consumed raw are probable sources of these microorganisms and attention should be paid to countries where sanitation is precarious (ALMEIDA FILHO, 2008).

Its main objective is to analyze the salads consumed raw by students, prepared in the cafeteria of the Vitória da Conquista campus, from a microbiological point of view.

MATERIALS AND METHODS

For the collection of salad samples, a plastic container with a lid was used, exclusively for this purpose, that is, to store the samples. They were then stored in a Styrofoam box containing an ice plate and immediately transported to the laboratory.

The analyzes were carried out by the server Dr. Roseane Machado, coordinator of the food and water laboratory, at the University of Southwest Bahia/UESB, Vitória da Conquista campus, This laboratory represents a reference for the municipality and region.

The methodology instituted in the UESB laboratory for the count of total and thermotolerant coliforms (450C) was the technique of the most probable number-MPN. For the serial decimal dilution of the samples, the first dilution (10⁻¹) of 225 ml of diluent (0.1% peptone water) was performed, plus 25g of the sample and homogenized. Then, 1 ml of the first dilution was transferred to 9 ml of the same diluent for the preparation of the second dilution (10⁻²), for the preparation of the third dilution (10⁻³), the same procedure was performed, transferring 1 ml of the second dilution to 9 ml of diluent, totaling three dilutions.

For the total coliform count, 1 ml of each dilution was added in test tubes containing 10 ml of Lauryl Sulfate Tryptose Broth (LST). The tubes were incubated in an incubator for 48 hours at 35°C, after which it was observed if there was gas production and turbidity of the tubes. From each tube with gas production, a loaded elevation of each culture was transferred to tubes containing Bright Green Bile 2% (VB) Broth and incubated in an incubator at 35°C for 48 hours. It was observed that BV tubes with gas production and turbidity, meaning the presence of total coliforms (APHA, 2015)

The count of thermotolerant coliforms was carried out from LST tubes with turbidity and gas production, where a loaded elevation of each culture was transferred to tubes with E. coli broth (EC). Then they were incubated in a water bath at 45°C for 24 hours, then the number of EC tubes with gas production was observed, confirming the growth of thermotolerant coliforms (APHA, 2015).



The results were analyzed based on the dilutions and the number of positive samples of the confirmatory test, then compared with the MPN table of the Bacteriological Analytical Handbook.

RESULTS

The results of the total coliform and *E. coli* analyses performed on the salads served to the students can be seen in Table 1. It is observed that the results obtained indicate that the 4 samples analyzed (100%) were outside the microbiological standards established in Normative Instruction No. 60/2019, since the sanitary conditions are unsatisfactory for human consumption, due to the analytical results being above the established limit. Regarding total coliforms, 4 (100%) of the samples were unhygienic Regarding *E. Coli* 3 (75%) they were well above the current established standard, which indicates that at some stage of the handling of the vegetables there was fecal contamination in the food, thus, the result indicates, with total safety, information about the hygienic conditions of the salads.

Data from the studies by PALÚ et al. (2002); PAULA et al. (2003); JUNIOR et al. (2012) are similar and corroborate their similarity with the results presented. They describe the presence of total coliforms and *E coli* in samples of lettuce, compound and mayonnaise salads. All of them state that, due to hygienic-sanitary failures throughout the preparation process, raw meals promote food insecurity. Research carried out in daycare centers and schools in other states also found.

Table 1. Microbiological analysis in raw salads served in a restaurant, at a Higher Education Institution in Vitória da Conquista (BA) 2023.

Samples	Total Coliforms NMP/g VMP2 - Normative Instruction No. 60/2019 Not referenced	Escherichia Coli NMP/g VMP2 - Normative Instruction N°60/2019 10
A	1100	210
B	3,6	Absent
C	150	93
D	150	93

Source: Survey data, 2023

FINAL CONSIDERATIONS

With the evaluations of the salads, even though it is a preliminary study, it was found that the restaurant of the higher education institution provided the students with raw salads with a high count of microorganisms such as total coliforms and *E. Coli*, which can compromise the health of the students. The results indicate contamination caused by the handlers, which can be deduced from the absence of hygienic-sanitary control of the production processes of raw



salads.

Thus, it is necessary for restaurant workers to be trained more frequently, to understand the importance of hand hygiene, to practice what is established in Good Handling Practices when preparing food, because it contains the real description of the technical procedures. Following all of these procedures prevents foodborne illnesses (DTAs).

It is expected that the PRPGI will provide public notices so that the research group Society: environment and health can continue this research, due to the importance of eating food within the microbiological patterns of food consumption.

Keywords: Student cafeteria, Raw salads, Microbiological analysis, Infectious diseases.



REFERENCES

- Nacional de Alimentação Escolar (PNAE). (2009b). Diário Oficial da União. Retrieved from http://educacaointegral.mec.gov.br/images/pdf/res_cd_38_16072009.pdf. Accessed August 18, 2023.
- CECANE PARANÁ. (2010). A agricultura familiar e o programa nacional de alimentação escolar – PNAE. Curitiba.
- Conceição, A. A. da. (2019). História da alimentação escolar no Brasil: algumas questões sobre políticas públicas educacionais, cultura escolar e cultura alimentar. In Anais do 30 Simpósio Nacional de História, Recife, July 15-19, 2019.
- Cunha, M. A. de. (2006). Métodos de detecção de microrganismos indicadores. Saúde & Ambiente em Revista, Duque de Caxias, 1(1), 9-13.
- Franco, B. D. G. M., & Landgraf, M. (2008). Microbiologia dos alimentos. São Paulo: Atheneu.
- Junior, J. P., Gontijo, E. E. L., & Silva, M. G. (2012). Perfil parasitológico e microbiológico de alfaces comercializadas em restaurantes self-service de Gurupi – TO. Revista Científica do ITPAC, 5(1), 1-8.
- Nascimento, M. S., & Catanozi, M. P. L. M. (2003). Avaliação microbiológica de frutas e hortaliças frescas, comercializadas no município de Campinas – SP. Rev Hig Alim, 17, 114-115.
- Oliveira, M. L. S., et al. (2006). Análise microbiológica de alface (*Lactuca sativa* L.) e tomate (*Solanum lycopersicon* L.) comercializados em feiras-livres da cidade de Belém, Pará. Higiene Alimentar, 19(143), 96-101.
- Paula, P., Rodrigues, P. S. S., Tortora, J. C. O., Uchôa, C. M. A., & Farage, S. (2003). Contaminação microbiológica e parasitológica em alfaces (*Lactuca sativa*) de restaurantes self-service, de Niterói, RJ. Rev Soc Bras Med Trop, 36, 535-537.
- Palú, Â. P., et al. (2002). Avaliação microbiológica de frutas e hortaliças frescas, servidas em restaurantes self service privados, da Universidade Federal do Rio de Janeiro. Higiene Alimentar, 16(100), 67-74.
- Takayanagui, O. M., et al. (2001). Fiscalização de verduras comercializadas no município de Ribeirão Preto, SP. Revista da Sociedade Brasileira de Medicina Tropical, 34(1), 37-41.