Chapter 24

Toxicological aspects and the sustainability factor in Brazilian beekeeping

Scrossref 💿 https://doi.org/10.56238/methofocusinterv1-024

Andréia Alves Rosa

Graduate Program in Agribusiness (PROPAGA), Darcy Ribeiro University Campus University of Brasília, Brasília, DF, 70910-900, Brazil E-mail: arrosalves@gmail.com

Artur Guerra Rosa

Graduate Program in Agribusiness (PROPAGA), Darcy Ribeiro University Campus University of Brasília, Brasília, DF, 70910-900, Brazil E-mail: rturguerra921@hotmail.com

ABSTRACT

Brazilian legislation conceptualizes honey as the "food product produced by honey bees, from the nectar of flowers or secretions from living parts of plants or from excretions of plant-sucking insects that are on living parts of plants, which bees collect, transform, combine with specific substances of their own, store and mature in the honeycombs. The term "sustainability" is derived from the word sustainable (latin "sus-tenere") which defines the ability to sustain or support certain conditions. Currently, it has been widely used in socioeconomic and environmental issues, mainly. Studying toxicological aspects in Brazilian beekeeping united with sustainability was the objective of this work.

Keywords: Beekeeping, Sustainability, Toxicological.

1 INTRODUCTION

According to Anacleto et al. (2009), the characteristics of honey, such as aroma, color and taste, depend both on the composition of the nectar of the plant species (flowering of geographical regions) and on the species of the bee that produces it, being the climate, soil and the form of management by the beekeeper factors that cause minor effects on its characteristics (FELSNER, 2001).

According to LIANDA (2009), the chemical composition of honey is defined as a complex mixture of carbohydrates, enzymes, acids, amino acids, minerals, aromatic substances, vitamins, pigments, waxes and pollen grains. And, more broadly, it can be said that honey is composed of water (17%), glycides (80%) and other substances (3%), such as enzymes [(invertase, amylase (diastasis) and glucose oxidase)], amino acids (the most abundant proline), organic acids, flavonoids and mineral matter.

According to Almeida-Muradian (2009), honey is the product of the best known beekeeping and marketed in the world, being all the products of the hive (honey, bee pollen, royal jelly, propolis, beeswax and apitoxin) registered with the Ministry of Agriculture, Livestock and Supply (MAPA).

China is the largest honey producer in the world, although the product does not have good quality, according to the Brazilian Confederation of Beekeeping (CBA, 2013). The United States is in second place, followed by Argentina in third. Brazil, according to SEBRAE (2012), is in eleventh place, and fifth, as an

exporter. In 2011, Brazil exported more than 20,000 tons of honey, representing an increase of more than 24% compared to 2010 in values, and an increase in net weight of more than 18%. However, in August 2012, when compared to the export of this product in the same period of 2011, there was a drop of about 70% in value and 68% in net weight.

This growth in Exports of Brazilian Honey, until 2011, occurred after the international embargo with the European Community, which occurred in the years 2006 to 2008, due to the allegation of noncompliance with sanitary requirements in relation to the control of pesticide residues and veterinary drugs (PAULA, 2008). From this event, there was a joint effort by beekeepers and institutions to produce quality food, free of antibiotics and providing food security (Al-WAILI et. Al. 2012). The decrease observed from August 2012 was due to the period of great drought in that period, followed by intense and heavy rains that washed the nectar of flowers, thus decreasing productivity (SEBRAE, 2013).

In Brazil, honey is mainly produced by Africanized bees that, due to its genetic characteristics, dispenses with the use of antibiotics and pesticides, such as acaricides and fungicides. In some Brazilian regions, according to embrapa meio-norte researchers (Rangel, 2011), honey is recognized worldwide as a food free of toxic products and, due to this worldwide recognition, honey produced in Brazil is sought worldwide. However, in other countries, Krupke et al (2012) and Blacquière et al (2012), presented studies showing the presence of agricultural pesticides present in bee populations, which in addition to compromising the health of hives and their productivity, the presence of this contaminant may also offer a health risk consumer.

2 DEVELOPMENT

2.1 BRAZILIAN LEGISLATION ON TRACE ELEMENTS IN HONEY

With the objective of defining safe levels in relation to the presence of toxic substances in samples of meare and other foods, the Ministry of Agriculture, Livestock and Supply (MAPA), through Normative Instruction No. 42 of December 20, 1999 (Brazil, 1999), issued the so-called National Waste Control Plan on Animal Products and, in May 2009, through Normative Instruction No. 14, the Program for The Control of Waste and Contaminants in Meat (Beef, Poultry, Pork and Equina), Milk, Honey, Eggs and Fish. In this last instruction, Annex IV refers to the Program for The Control of Residues and Contaminants in Honey (PCRN)(BRASIL, 2009).

Table 1 shows the maximum concentration of some metals in honey, allowed by Brazilian legislation, according to Ribeiro (2010).

|--|

| | MAXIMUM PERMITTED CONCENTRATION (µg.g-1) | | | | | | |
|---------------|--|-------------|--------------|---------|--|--|--|
| Elements | Dec. 55871/65 | Port. 11/87 | Port. 685/98 | IN 9/14 | | | |
| Arsenic (As) | 1,0 | ND | 1,0 | 0,5 | | | |
| Cadmium (Cd) | 1,0 | ND | ND | 0,5 | | | |
| Lead (Pb) | 0,8 | ND | ND | 0,5 | | | |
| Copper (Cu) | 30,0 | ND | 10,0 | ND | | | |
| Chromium (Cr) | 0,1 | 0,1 | ND | ND | | | |
| Mercury (Hg) | 0,01 | ND | ND | 0,5 | | | |
| Nickel (Ni) | 5,0 | ND | ND | ND | | | |
| Selenium (If) | 0,05 | ND | ND | ND | | | |
| Zinc (Zn) | 50 | ND | ND | ND | | | |

Dec. 55871/65 - Decree No. 55871/65 (BRASIL, 1965)

Port. 11/87 - Ordinance No[.] 11/87 (BRASIL, 1987) Port. 685/98 - Ordinance No. 685/98 (BRASIL, 1998) IN 14/09 - Normative Instruction No. 14/09 (BRASIL, 2009) ND = Not defined by Legislation

Source: Ribeiro (2010) - Modified

In addition to traces of heavy metals, the PNCR also cites other contaminants, among them , antibiotics (Penicillin, Chloramphenicol, Streptominica, Tetracycline, Erythromycin, Neomycin, Oxytetracycline, Chlortetracycline); Sulfonamides (Sulfadimetoxin, Sulfametazin, Sulfatiazol, in addition to other drugs (Nitrofurazone and Furazolidone)(BRASIL, 2007).

In Table 2, Pittella (2009) demonstrated the maximum limits of antibiotic residues allowed in honey, according to the Brazilian Legislation, as well as the number of trials performed and the techniques used in the study conducted in 2009.

| Table 2: Maximum Limits of Antibiotic Residues Allowed by Brazilian Legislation. | | | | | | |
|--|-----------------------|-----------|----------|-----------|--------------------|--|
| Group | Analyt | Technique | LQ | LMR/NA* | ^{No} Test | |
| | | | (µgKg/L) | (µg/kg/L) | Items | |
| | Chlortetracycline (a) | | | | | |
| | Oxytetracycline (a) | HPLC- | 25 | 200 | 60 | |
| | Tetracycline (a) | UV | | | | |
| | Sulfatiazole (b) | | | | | |
| | Sulfametazin (b) | HPLC- | 25 | 100 | 60 | |
| Antimicrobials | Sulfadim etoxin | UV | | | | |
| | (b) | | | | | |
| | Sulfadiazine | | | | | |
| | Sulfaquinoxaline | | | | | |
| | Sulfaclorpyridazine | | | | | |
| | Nitrofurazone | LC/ MS- | 0,5 | 1(1)* | 30 | |
| | Furazolidona | MS | | | | |
| | Furaltadona | | | | | |
| | Nitrofurantoin | | | | | |
| | Tylosin | LC/ MS- | 0,5 | 10 | 10 | |
| | | MS | | | | |

(a) MRL refers to the sum of all tetracycline

(b) The MRL refers to the sum of all Sulfonamides (*) NA - Level of action

HPLC - UV - Liquid alpha efficiency chromatography and visible ultraviolet detector LC/MS-MS - Liquid chromatography coupled to tandem mass spectrometry Source: Normative Instruction No. 9 of March 30, 2007. MAP. Pittella (2009) - Modified

This same Normative Instruction No. 9 of the Map also includes pesticide residues, such as another class of contaminants in medes and provides, according to Pittella (2009), the execution of analyses for the detection of these contaminants as well as the Maximum Residue Limit (MRL) for various types of halogenated, organochlorinated compounds, carbamates, pyrethroides and organophosphates in mes.

In Table 3, Pittella (2009) demonstrates the National Honey Waste Control Plan (PNCR-2007) containing, in addition to the number of tests performed and the techniques used, the Maximum Residue (MRL) and Quantification Limits (QS), recommended by the IN in question.

 Table 3: Limits Maximum from Waste (MRL) e Limits from Quantification (LQ) from Waste from Pesticides Permitted by law Brazilian.

| Group | Analyt | Matrix | Technique | LQ | LMR/NA* | In Items |
|-------------------|--------------|--------|-----------|-----------|--------------|----------|
| | | | | (µg/kg OR | (µg/kg or | test |
| | | | | μg/L) | μg/L) | |
| | Aldrin | | | 10 | 10 | |
| | Alpha- | | | 10 | 10 | |
| | Endosulfan | | | | | |
| | 4.4-DDE | | | 10 | 10 | |
| Compounds | 4.4-DDD | | | 10 | 10 | |
| Halogenated And | 4.4- DDT | | | 10 | 10 | |
| Organochlorinated | | | | | . | |
| | Dodecachlor | Honey | CG/EM | 10 | 10 | 15 |
| | Endrin | | | 10 | 10 | |
| | Iprodione | | | 50 | 50 | |
| | Tetradifone | | | 20 | 20 | |
| | Captana | | | 50 | 50 | |
| | Liudane | | | 10 | 10 | |
| | Vinclozolin | | | 20 | 20 | |
| | Permethrin | | | 64,6 | 64,6 | |
| | Cyfluthrin | | | 44 | 44 | |
| | Fenpropatrin | | | 10 | 10 | |
| Carbamates And | Deltamethrin | Honey | CG/EM | 33,8 | 33,8 | 15 |
| Pyrethroides | Carbofuran | | | 82 | 82 | |
| | Carbaril | | | 37 | 37 | |
| | Pyriphos | | | 36 | 36 | |
| | Methyl | | | | | |
| | Chlorpyrifos | Honey | CG/EM | 24,3 | 24,3 | 30 |
| Organophosphates | Dimetoate | | | 20 | 20 | |
| | Dissulfoton | | | 10 | 10 | |
| | Paration | | | 36,9 | 36,9 | |
| | Fenamifós | | | 17,1 | 17,1 | |
| | Terbuphos | | | 13 | 13 | |
| | Profenofós | | | 36 | 36 | |

LQ - LMR quantification limit - Maximum residue limit (*) NA - Level of Action

CG-MS – Gas chromatography coupled to mass spectrometry Source: Normative Instruction No. 9 of March 30, 2008. MAP Source: Pittella (2009) - Modified

2.2 ANALYSIS OF RESULTS

Sustainability is a well-known word in agriculture since the 19th century, but has been gaining prominence since the 1980s and 1990s with the issue of social concern in relation to the environmental issue.

According to Silva et al (2008), since the Stockholm Conference in 1972 society began to worry also about the socio-environmental issue in issues related to world development versus widespread degradation of the environment and natural resources. However, even after all the moves that have occurred around the concern of sustainable practice such as the creation of the World Commission on Environment and Development (CMMAD) in 1983 by the United Nations (UN), and the United Nations Conference on Environment and Development in Rio de Janeiro, in 1992, it is observed that the attention of managers of the numerous associations created with the objective of making sustainability, both in the beekeeping sector and in other sectors, an emerging factor capable of integrating economic growth, with social development and environmental protection, still remains much more focused on the economic issue than in social and environmental environment.

Silva et al (2008), explains in their study that, although beekeepers have individual concern so far in relation to environmental and social issues (e.g., afforestation of beekeepers' properties, donation of honey plants to schools, preservation and environmental awareness , the donation of beeproducts to entities, as well as transmitting to society the benefits of bee products for public health in addition to the importance of bees in the environmental context), they become associated, in the most often , the various class associations of the beekeeping sector with the intention of acquiring space and greater profit in the economic market.

3 FINAL CONSIDERATIONS

Beekeeping is, according to Both et al (2009), one of the only activities, among the numerous existing production processes, which presents better socio-economic and environmental conditions for the sustainability process, because it generates employment and consequently income for producers (beekeepers); decreases the rural exodus and, beekeeping production is linked to the regeneration of natural vegetation (withdrawal from nectar, pollination etc). However, with the progress of agricultural processes (agriculture, livestock, deforestation, among others) there is a great fall in bee production, since, in most cases observed, beekeepers have great difficulties for the development of beekeeping, such as the lack of organization and programs for the development of producers; lack of information on the use of veterinary antibiotics (e.g. pesticides among other products, which can lead to both the death of honey bees and contamination of the bee product.

REFERENCES

ALMEIDA-MURADIAN, L. B. Qualidade dos produtos apícolas e otimização quimiométrica dos métodos de análise do mel por espectroscopia no infravermelho (FT-IR ATR). Tese de Livre-Docência da FCF/USP, 2009.

AL-WAILI, N, et al. Antibiotic, Pesticide, and Microbial Contaminants of Honey: Human Health Hazards. The Scientific World Journal, Volume 2012, Article ID 930849, 9 p.

ANACLETO, D. A.; SOUZA, B. A.; MARCHINI, L. C.; MORETI, A. C. C. C. Composição de amostras de mel de abelha Jataí (Tetragonisca angustula latreille, 1811). Ciênc. Tecnol. Aliment., Campinas, 29(3): 535-541, jul.-set. 2009.

BLACQUIÈRE, T, et al. Neonicotinoids in bees: a review on concentrations, side- effects and risk assessment. Ecotoxicology, 21, p. 973–992, 2012.

BOTH, J. P. C. L., KATO, O. R., OLIVEIRA, T. F. Perfil socioeconômico e tecnológico da apicultura no município de Capitão Poço, Estado do Pará, Brasil. Amazônia: Ci. & Desenv., Belém, v. 5, n. 9, jul./dez. 2009.

BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa 11 de 20 de outubro de 2000, Regulamento Técnico de identidade e qualidade do mel. Disponível em: http://extranet.agricultura.gov.br/sislegis-consulta/ consultar Legislacao. do?operacao=visualizar&id=7797. Acesso em: 05/06/2021.

BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa nº 9 de 30 de março de 2007. Programas de Controle de Resíduos e Contaminantes em Carne (Bovina, Aves, Suína e Eqüina), Leite, Mel, Ovos e Pescado do exercício de 2007. Diário Oficial da União Brasília 04/04/2007 Seção 1, p. 7, 2007.

BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa nº 14, de 25 de maio de 2009 – Aprova os Programas de Controle de Resíduos e Contaminantes em Carnes (Bovina, Aves, Suína e Equina), Leite, Mel, Ovos e Pescado para o exercício de 2009 e seu anexo: "Programa de Controle de Resíduos e Contaminantes em Mel – PNCRC/2009". Diário Oficial da União, Brasília, DF, 2009.

BRASIL. Regulamento técnico Mercosul: Identidade e qualidade do mel. Resolução GMC nº15/94. Montevidéu, 1999. Disponível em: http://www.extranet.agricultura. gov. br/consultasislegis/do/consultaLei?op=viewTextual&código=6020. Acesso em: 09/06/2021.

CBA – Confederação Brasileira de Apicultura. Qualidade do mel brasileiro impulsiona crescimento do setor. Disponível em: http://www.brasilapicola.com.br/mercado-e-cia-05012012-qualidade-do-mel-brasileiro-impulsiona-crescimento-do-setor. Acesso em: 05/06/2021.

FELSNER, M. L. Tese de Doutorado. Caracterização de méis monoflorais de eucalipto e de laranja do Estado de São Paulo por técnicas termoanalíticas. Universidade de São Paulo-USP, São Paulo, 2001.

KRUPKE, C. H., et al. Multiple Routes of Pesticide Exposure for Honey Bees Living Near Agricultural Fields. PLoS ONE V. 7, N.1, e29268, 2012.

LIANDA, R. L. P. Tese de Doutorado. Perfil de substâncias fenólicas de méis brasileiros por cromatografia líquida de alta eficiência e avaliação do potencial antioxidante. Universidade Federal Rural

Methodology focused on the area of interdisciplinarity: Teenager with leprosy and self-stigma: The role of education do Rio de Janeiro. Rio de Janeiro, 2009. PAULA, J. Mel de Brasil, as exportações brasileiras do mel no período 2000/2006 e o papel do SEBRAE, Brasília, DF. SEBRAE 2008.

PITTELLA, M. C. Determinação de resíduos de agrotóxicos em mel de abelhas (Apis sp) por cromatografia gasosa acoplada à espectrometria de massas. Dissertação de Mestrado. Universidade Federal de Minas Gerais. Belo Horizonte, MG, 2009.

RANGEL, R. Apicultura: Mel brasileiro conquista o mercado externo. Revista Inovação em Pauta (FINEP), 10^a edição – Nov/dez 2010-jan/2011. Disponível em: http://www.finep.gov.br/imprensa/noticia.asp?cod_noticia=2485. Acesso em 03/06/2021.

RIBEIRO, R. O. R. Elementos traço em méis de abelha (Apis mellifera) do estado do Rio de Janeiro, Brasil: Influências da Sazonalidade. Dissertação Mestrado. Universidade Federal Fluminense, Niterói, RJ. 2010.

SEBRAE. Serviço Brasileiro de Apoio às Micro e Pequenas Empresas. Agência Sebrae de Notícias. Aumento das exportações amplia mercado do mel brasileiro. Disponível em: http://www.agenciasebrae.com.br/noticia/12816861/agronegocios/aumento-das- exportacoes-ampliamercado-do-mel-brasileiro/. Acesso em 14/06/2021.

SEBRAE. Serviço Brasileiro de Apoio às Micro e Pequenas Empresas. Agência Sebrae de Notícias. Aumento das exportações amplia mercado do mel brasileiro. Disponível em: http://www.sebrae.com.br/setor/apicultura/sobre- apicultura/mercado/exportacoes. Acesso em 10/07/2021.

SILVA, T. N.; LENGLER, L. Sustentabilidade, Empreendedorismo e Cooperação em Associações de Apicultores da Região Central do Rio Grande do Sul. Revista da Micro e Pequena Empresa, Campo Limpo Paulista, v.2, n.2, p.40-57, 2008.