 <https://doi.org/10.56238/alookdevelopv1-052>

Jefferson de Souza Busso

Institute of Biosciences, Letters and Exact Sciences (IBILCE) – São Paulo State University “Júlio de Mesquita Filho” (UNESP) – campus of São José do Rio Preto/ SP/ Brazil.

Multiuser center Biomolecular, Physics Department, Instituto de Biociências, Letras e Ciências Exatas (IBILCE) – Universidade Estadual Paulista “Júlio de Mesquita Filho” (UNESP) – campus de São José do Rio Preto/ SP/ Brazil.

Orcid: <https://orcid.org/0000-0002-6567-3367>

Jéssica Maróstica de Sá

Institute of Biosciences, Letters and Exact Sciences (IBILCE) – São Paulo State University “Júlio de Mesquita Filho” (UNESP) – campus of São José do Rio Preto/ SP/ Brazil.

Multiuser center Biomolecular, Physics Department, Instituto de Biociências, Letras e Ciências Exatas (IBILCE) – Universidade Estadual Paulista “Júlio de Mesquita Filho” (UNESP) – campus de São José do Rio Preto/ SP/ Brazil.

Orcid: <https://orcid.org/0000-0003-0025-8311>

Marcelo Andres Fossey

Institute of Biosciences, Letters and Exact Sciences (IBILCE) – São Paulo State University “Júlio de Mesquita Filho” (UNESP) – campus of São José do Rio Preto/ SP/ Brazil

Multiuser center Biomolecular, Physics Department, Instituto de Biociências, Letras e Ciências Exatas (IBILCE) – Universidade Estadual Paulista “Júlio de Mesquita Filho” (UNESP) – campus de São José do Rio Preto/ SP/ Brazil

Orcid: <https://orcid.org/0000-0002-2012-388X>

Ícaro Putinhon Caruso

Institute of Biosciences, Letters and Exact Sciences (IBILCE) – São Paulo State University “Júlio de Mesquita Filho” (UNESP) – campus of São José do Rio Preto/ SP/ Brazil.

Multiuser center Biomolecular, Physics Department, Instituto de Biociências, Letras e Ciências Exatas (IBILCE) – Universidade Estadual Paulista “Júlio de Mesquita Filho” (UNESP) – campus de São José do Rio Preto/ SP/ Brazil.

Orcid: <https://orcid.org/0000-0003-4464-0520>

Fátima Pereira de Souza

Institute of Biosciences, Letters and Exact Sciences (IBILCE) – São Paulo State University “Júlio de

Mesquita Filho” (UNESP) – campus of São José do Rio Preto/ SP/ Brazil

Multiuser center Biomolecular, Physics Department, Instituto de Biociências, Letras e Ciências Exatas (IBILCE) – Universidade Estadual Paulista “Júlio de Mesquita Filho” (UNESP) – campus de São José do Rio Preto/ SP/ Brazil

Orcid: <https://orcid.org/0000-0002-4731-4977>

E-mail: fatima.p.souza@unesp.br

ABSTRACT

Intestinal microbiota plays an important role in the human organism homeostasis and it is subject to several perturbations caused by diet and environmental factors, inducing dysbiosis. The search for compounds can help rebuild the intestinal microbiota because sometimes this is the only treatment option for intestinal dysbiosis. Phenolic compounds are neutralizing free radicals, showing a mutual relationship with foods rich in flavonoids, so they might be helpful during the restoration of the intestinal microbiota. This study analyzed the direct effect of these compounds on *Lactobacillus rhamnosus*, cultivated with and without flavonoids in MRS broth, in different growth phases and concentrations. After that, were exposed to enzymes that are involved in the human digestive process to analyze any alteration. Flavonoids quercetin, rutin, hesperetin, and hesperidin were individually tested. The results showed that the presence of flavonoids increased the number of colony-forming units (CFU) compared to the control group (without flavonoids), especially when added before the lag phase. In the digestibility test, quercetin showed a minimal effect for the acidic stomach stage, but for the other steps mimicking the further digestive system, there was no difference compared to the control, all being killed after bile salts. Therefore, we concluded flavonoids are helpful in the proliferation of the tested stain; however, it is necessary to develop more efficient methods so that the flavonoids can overcome the digestive system and reach the intestine with total efficiency.

Keywords: Flavonoids, probiotics, prebiotics, dysbiosis, microbiology, molecular biology.

1 INTRODUCTION

The human body is colonized by several microorganisms during life, including bacteria, archaea, viruses, and eukaryotes, in a complex ecosystem that coexists with its host in a mutualistic way. These microorganisms settle on body surfaces, such as skin and urogenital, respiratory, and gastrointestinal tracts [1,2]. The gastrointestinal tract, especially the small intestine, stands out with conditions for the proliferation of these organisms, called probiotics, constituting intestinal microbiota [3]. Probiotics are defined as live microorganisms that when administered properly and consumed safely, confer benefits to the health of their host and studies show the species to use for treatments [4,5]. Azad and collaborators revealed that the *Bifidobacterium* and *Lactobacillus* genera have the largest number of species, being the most used in research [6]. The small intestine provides nutrients and favorable conditions for probiotics proliferation, such as temperature and anaerobiosis. In turn, intestinal microbiota metabolism assists the fermentation of carbohydrates, vitamin synthesis, increasing intestinal permeability, and epithelial defense mechanism to form the mucosal barrier [7,8].

Intestinal microbiota health is intrinsically related to the host health, as shown in germ-free animal studies, that the absence of these microorganisms made them more susceptible to the development of diseases, such as obesity, type II diabetes, hypertension, and inflammatory bowel diseases [9-13]. Scheperjans and coauthors observed that the abundance of the *Prevotellaceae* family has a protective effect on human Parkinson's disease, while its decrease may be a biomarker for this disease [14]. The maintenance of microbiota is essential for host homeostasis, which can contribute during the treatment of several diseases. Compounds that help probiotics proliferate in a safe and lasting way may be the key to restoring the altered microbiota, in addition to being able to optimize foods that supplement the probiotics [15,16].

Conjugation between probiotics and flavonoids was described as mutualistic in the review by Cardona and coauthors, who reported that some flavonoids are absorbed in the small intestine after being metabolized by specific bacteria, and on the other hand, the intake of foods rich in flavonoids modulates probiotic populations, contributing to their maintenance, stimulating the growth of beneficial bacteria, inhibiting the proliferation of pathogenic bacteria [17]. Flavonoids are secondary metabolites of plants, found in leaves, fruits, and seeds, present in the human diet. Its structure has two benzene rings linked to a pyran ring, rich in hydroxyls and electron donor, which confers antioxidant activities, widely studied, by reacting and inactivating superoxide anions, singlet oxygen, lipid peroxide radicals and/or stabilizing free radicals involved in the oxidative process through hydrogenation or complexation with oxidizing species [18,19]. Biologically, the diversity of known molecules plays an important hepatoprotective, antibacterial, anti-inflammatory, anticancer, and antiviral role [20].

Several studies investigated the relationship between foods rich in polyphenols and probiotics, such as cocoa derivatives, grape derivatives, red pitaya pulp, and berries, which have shown significant

promise in optimizing probiotics, functioning as prebiotics [21-24]. Given the evidence presented, the combination of flavonoids and probiotics shows promise and can expand knowledge of how this complex ecosystem works and interacts with the host. In addition, these studies show evidence of how the body can reestablish homeostasis when subjected to changes. This study tested the prebiotic effect of these molecules on the isolated condition of the probiotic present in the human intestine microbiota and its effect on simulated gastrointestinal digestion.

2 MATERIALS AND METHODS

2.1 STAIN AND INOCULUM PREPARATION

The strain was cultivated in MRS broth (Man, Rogosa & Sharpe, from Kasvi, São José dos Pinhais, PR, Brazil) at 37 °C overnight (14-16 hours) until reach stationary phase, with DO 1.8-2.0, previously determinate in this study and compared with the literature [25]. All tests and controls were serially diluted in saline solution 0,9% (0,9 g/mL NaCl), plated on MRS Agar (from Kasvi, São José dos Pinhais, PR, Brazil), and grew anaerobically for 24 h at 37 °C. After incubation time, bacterial colonies were counted and compared with the control, expressed in percentage. Error percentage was calculated through the ratio between the means and standard deviation of the control and tests.

2.2 MOLECULE CHARACTERISTICS AND EFFECTS ON PROBIOTIC GROWTH


Flavonoids used in this study were Quercetin (QUE), Rutin (RUT), Hesperidin (HET), and Hesperidin (HED), all purchased from Sigma-Aldrich, St. Louis, MO, United States, with $\geq 95\%$ purity. RUT is derived from QUE, as well as HED derived from HET, with both molecules increased with carbohydrates rhamnose and glucose (Figure 1). The stock solutions of 5 mM were prepared in 99,9% DMSO (Dimethyl sulfoxide, from Sigma-Aldrich, St. Louis, MO, United States) using the molar absorption coefficients of 21.880 M⁻¹ cm⁻¹ (375 nm), 18.800 M⁻¹ cm⁻¹ (358 nm), 33.232 M⁻¹ cm⁻¹ (290 nm) and 18.170 M⁻¹ cm⁻¹ (295 nm), respectively, and determinate by UV-Vis Spectrophotometer (BioMate™ 3S, ThermoFisher, USA), equipped with 1.0 cm quard cells of path length, on the range of 250-500 nm absorption spectra.

DMSO was tested separately as solvent control at the range of 0.1% and 1.0% on absolute concentration (99,9 %), and flavonoids were tested on low (10 µM) and medium (50 µM) concentrations. Solvent and phenolic compounds were added in MRS broth for 2 hours on the stationary phase, after growth for 16 hours anaerobically, for analyzing the effect between the concentrations. The medium concentration was added before the lag phase and allowed growth for 16 hours for analyzing the influence during proliferation.

2.3 PRELIMINARY SIMULATION OF GASTROINTESTINAL DIGESTION

The gastrointestinal simulation process was carried out in stages, using digestive enzymes (Sigma, St. Louis MO, USA), under constant agitation. The pH of each solution stage was adjusted using 1M HCl or 1M NaHCO₃. To simulate mastication, 1.2 mL/min of salivary solution (0.33 mg of α -amylase/mL of 1 mM CaCl₂) was added at neutral pH of 6.9, under intense agitation (200 rpm), for 2 minutes. For the esophagus-stomach step, stirring at 130 rpm and pepsin solution (25 mg/mL of 0.1 M HCl) was added at a rate of 0.05 mL/mL of inoculum and pH 2.0 for 90 min. The intestinal solution was prepared with 2 g/L of pancreatin and 12 g/L of bile salts, diluted in a 0.1 M NaHCO₃ solution, stirred at 45 rpm, with pH 5.0 for 30 min. Finally, the pH was adjusted to 6.5 with NaHCO₃ to simulate the ileal portion for 60 min [26,27]. After each step, a 100 μ L-aliquot was serially diluted in saline solution and plated on MRS agar for 24 hours.

Figure 1. Chemical structure of (A) hesperetin, (B) quercetin, (C) hesperidin, and (D) rutin.



3 RESULTS

The first test was with DMSO, added at a rate of 0.1% (1 μ L/mL of MRS broth) and 1.0% (10 μ L/mL of MRS broth) for 2 hours on the stationary phase, after growth overnight. The percentage of growth increased to 32% and 69%, respectively. When 1.0% was added before the lag phase and growth anaerobically overnight, the CFU was increased to 74%.

Phenolic compounds used in this study also increased the CFU, overcoming the DMSO effect. The difference between low and medium concentration was not so significant when added in the stationary phase after growing overnight, resulting, respectively, in an increase of 60% and 78% for quercetin, 98%, and 106% for rutin, 57% and 78% for hesperetin, and 126% and 134% for hesperidin.

However, when medium concentrations were added before the lag phase and growth overnight, large increases occurred compared to the control, with hesperidin having the greatest increase (318%), followed by

rutin (237%), hesperetin (131%), and quercetin (101%). All percentages are expressed and compared in figure 2.

The preliminary simulation of gastrointestinal digestion was done with quercetin. The results showed that the molecule increased the formation of colonies in the stages of the mouth and stomach, and in the most acidic step, the stomach, its percentage increased by 426% (Figure 3).

4 DISCUSSIONS

DMSO is a widely used solvent due to its efficient physicochemical properties in solubilizing water-insoluble compounds, such as flavonoids. However, a high concentration of this solvent is toxic and may kill cells [28]. Then the solvent was tested as a control, and the results show that DMSO increased CFU in three treatments. The increase in CFUs may be a result of the strain's ability to metabolize the solvent and use it as a carbon source, as explained by Vdihya & Thatheyus in their study with DMSO and *Bacillus subtilis* [29]. Hesperidin and rutin had the highest increase, which might be related to the carbohydrates in the molecules, also being metabolized by bacteria as a source of carbon.

Figure 2. Effect of the presence of flavonoids on colony-forming units. Relation between the low and medium concentration on the percentual when added during the stationary phase. When the flavonoids were added before the lag phase, at medium concentration the percentual was intensely increased.





Figure 3. Simulation of gastrointestinal digestion. Comparison strain treated with quercetin and a significant increase in colony-formed units, during the control remained constant until the addition of bile salts in the duodenal stage.



Studies have shown how foods rich with flavonoids, this study has notable effects on the intestinal microbiota, such as quercetin, which stands out with its protective action against intestinal dysregulation during the use of antibiotics, increased intestinal permeability associated with the production of acids short-chain fatty acids, in addition to acting against pathogens, promoting balance in the microbiome [30-32]. Hesperidin proved to be effective in increasing populations of *Bifidobacterium* by influencing the action of IgA in mice [33]. Rutin action depends on how it's metabolized, producing different types of phenolic acids that can help in the bioavailability of other compounds [34], consolidating its prebiotic potential for these microorganisms and other benefits to the host.

The increase during the stomach stage might be related to the molecule propriety of the electron donor, which confers antioxidant activities [35]. This stage at control confirmed the tolerance of *L. rhamnosus* to acid pH [36,37], keeping colony-forming units close to 100%.

However, *L. rhamnosus* is also described as bile tolerant, and bile salts killed all cells in the duodenum stage. This result might be from its biological function in the digestion of lipids, damaging cell membranes and restricting the growth of some species of bacteria [38].

5 CONCLUSIONS

In conclusion, flavonoids presented an effect on the proliferation of the *L. rhamnosus* strain. This effect is even more potentiated when added medium before the proliferative phase, reaffirming the prebiotic effect of these compounds. The combination of probiotics and flavonoids is promising and can expand knowledge about how the microbiota works and your applications in several diseases

of the intestinal tract. Preliminary tests showed that these molecules might amplify proliferation in acid pH, but would not protect from all digestion stages.

REFERENCES

1. Sekirov, I.; Russell, S.L.; Antunes, L.C.; Finlay, B.B. Gut microbiota in health and disease. *Physiol Rev.* 2010;90(3):859-904. doi:10.1152/physrev.00045.2009
2. Morelli, L. Postnatal development of intestinal microflora as influenced by infant nutrition. *J Nutr.* 2008;138(9):1791S-1795S. doi:10.1093/jn/138.9.1791S
3. Ley, R.E.; Peterson, D.A.; Gordon, J.I. Ecological and evolutionary forces shaping microbial diversity in the human intestine. *Cell.* 2006;124(4):837-848. doi:10.1016/j.cell.2006.02.017
4. Guarner, F.; Schaafsma, G.J. *Probiotics. Int J Food Microbiol.* 1998;39(3):237-238. doi:10.1016/s0168-1605(97)00136-0
5. FAO/WHO. Guidelines for the Evaluation of Probiotics in Food. Food and Agriculture Organization of the United Nations/World Health Organization, London, Ontario, 2002.
6. Azad, M.A.K.; Sarker, M; Li, T.; Yin, J. Probiotic Species in the Modulation of Gut Microbiota: An Overview. *Biomed Res Int.* 2018;2018:9478630. doi:10.1155/2018/9478630
7. Berg, D.; Clemente, J.C.; Colombel, J.F. Can inflammatory bowel disease be permanently treated with short-term interventions on the microbiome? *Expert Rev Gastroenterol Hepatol.* 2015;9(6):781-795. doi:10.1586/17474124.2015.101303
8. Shi, N.; Li, N.; Duan, X.; Niu, H. Interaction between the gut microbiome and mucosal immune system. *Mil Med Res.* 2017; 4:14. doi:10.1186/s40779-017-0122-9
9. Turnbaugh, P.J.; Ley, R.E.; Mahowald, M.A.; Magrini, V.; Mardis, E.R.; Gordon, J.I. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature.* 2006;444(7122):1027-1031. doi:10.1038/nature05414
10. Frank, D. N.; St Amand, A.L.; Feldman, R.A.; Boedeker, E.C.; Harpaz, N.; Pace, N.R. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci U S A.* 2007;104(34):13780-13785. doi:10.1073/pnas.0706625104
11. Qin, J.; Li, Y.; Cai, Z.; et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature.* 2012;490(7418):55-60. doi:10.1038/nature11450
12. Yang, T.; Santisteban, M.M.; Rodriguez, V.; et al. Gut dysbiosis is linked to hypertension. *Hypertension.* 2015;65(6):1331-1340. doi:10.1161/HYPERTENSIONAHA.115.05315
13. Shi, N.; Li, N.; Duan, X.; Niu, H. Interaction between the gut microbiome and mucosal immune system. *Mil Med Res.* 2017;4:14. doi:10.1186/s40779-017-0122-9
14. Scheperjans, F.; Aho, V.; Pereira, P.A.; et al. Gut microbiota are related to Parkinson's disease and clinical phenotype. *Mov Disord.* 2015;30(3):350-358. doi:10.1002/mds.26069
15. Doodoo, C.C.; Wang, J.; Basit, A.W.; Stapleton, P.; Gaisford, S. Targeted delivery of probiotics to enhance gastrointestinal stability and intestinal colonisation. *Int J Pharm.* 2017;530(1-2):224-229. doi: 10.1016/j.ijpharm.2017.07.068
16. Rabêlo, C.A.C.; Patricio, M.F.B.P.; Naves, G.L.; Sullara, B.V.; Azevedo, H.C.S.S. Quantificação da microbiota presente em produtos lácteos industrializados comercializados como probióticos. *RECIMA21-Revista Científica Multidisciplinar - ISSN 2022;2675-6218, 3(5), e351418.*

doi:10.47820/recima21.v3i5.1418

17. Cardona, F.; Andrés-Lacueva, C.; Tulipani, S.; Tinahones, F.J.; Queipo-Ortuño, M.I. Benefits of polyphenols on gut microbiota and implications in human health. *J Nutr Biochem*. 2013;24(8):1415-1422. doi: 10.1016/j.jnutbio.2013.05.001
18. Birt, D.F.; Hendrich, S.; Wang, W. Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacol Ther*. 2001;90(2-3):157-177. doi:10.1016/s0163-7258(01)00137-1
19. Tafuri, N. F. Antiviral activity of plants extracts and flavonoids against Bovine herpesvirus 1 (BoHV1). Master's Dissertation in Biochemistry and Molecular Biology of Plants, Universidade Federal de Viçosa, Viçosa, 2011.
20. Kumar, S.; Pandey, A.K. Chemistry and biological activities of flavonoids: an overview. *ScientificWorldJournal*. 2013; 2013:162750. Published 2013 Dec 29. doi:10.1155/2013/162750
21. Succi, M.; Tremonte, P.; Coppola, R. Pre-cultivation with Selected Prebiotics Enhances the Survival and the Stress Response of *Lactobacillus rhamnosus* Strains in Simulated Gastrointestinal Transit. *Front Microbiol*. 2017; 8:1067. Published 2017 Jun 14. doi:10.3389/fmicb.2017.01067
22. Campanella, D.; Rizzello, C.G.; Fasciano, C.; et al. Exploitation of grape marc as functional substrate for lactic acid bacteria and bifidobacteria growth and enhanced antioxidant activity. *Food Microbiol*. 2017;65:25-35. doi:10.1016/j.fm.2017.01.019
23. Morais, S.G.G.; da Silva, G.C.B.; Dos Santos, L.M.; Martín-Belloso, O.; Magnani, M. Effects of probiotics on the content and bioaccessibility of phenolic compounds in red pitaya pulp. *Food Res Int*. 2019; 126:108681. doi: 10.1016/j.foodres.2019.108681
24. Pap, N.; et al. Berry polyphenols and human health: evidence of antioxidant, anti-inflammatory, microbiota modulation, and cell-protecting effects. *Current opinion in food science*. 2021;42:167-186. doi:10.1016/j.cofs.2021.06.003
25. Broeckx, G.; Kiekens, S.; Jokicevic, K.; Byl, E.; Henkens, T.; Vandenheuvel, D.; Lebeer, S.; Kiekens, F. Effects of initial cell concentration, growth phase, and process parameters on the viability of *Lactobacillus rhamnosus* GG after spray drying. *Drying Technology*, 2020;38:11, 1474-1492, doi: 10.1080/07373937.2019.1648290
26. ROLIM, F.R.L. Avaliação do Efeito Protetor de Queijo de Coalho Caprino na Sobrevivência de uma Nova Cepa com Potencial Probiótico. Dissertação de mestrado Universidade Federal da Paraíba, Paraíba, 2015.
27. Dos Santos, A.S.; de Albuquerque, T.M.R.; de Brito Alves, J.L.; de Souza, E.L. Effects of Quercetin and Resveratrol on in vitro Properties Related to the Functionality of Potentially Probiotic *Lactobacillus* Strains. *Front Microbiol*. 2019; 10:2229. doi:10.3389/fmicb.2019.02229
28. Santos, N.C.; Figueira-Coelho, J.; Martins-Silva, J.; Saldanha, C. Multidisciplinary utilization of dimethyl sulfoxide: pharmacological, cellular, and molecular aspects. *Biochem Pharmacol*. 2003;65(7):1035-1041. doi:10.1016/s0006-2952(03)00002-9
29. Vidhya, R.; Thatheyus, A.J. Biodegradation of Dimethylformamide Using *Bacillus subtilis*. *American Journal of Microbiological Research*, 2013; 1, 10-15. doi:10.12691/ajmr-1-1-3

30. Shabbir, U.; Rubab, M.; Daliri, E.B.; Chelliah, R.; Javed, A.; Oh, D.H. Curcumin, Quercetin, Catechins and Metabolic Diseases: The Role of Gut Microbiota. *Nutrients*. 2021;13(1):206. doi:10.3390/nu13010206
31. Shi, T.; Bian, X.; Yao, Z.; Wang, Y.; Gao, W.; Guo, C. Quercetin improves gut dysbiosis in antibiotic-treated mice. *Food Funct*. 2020;11, 8003–8013.
32. Ju, S.; Ge, Y.; Li, P.; et al. Dietary quercetin ameliorates experimental colitis in mouse by remodeling the function of colonic macrophages via a heme oxygenase-1-dependent pathway. *Cell Cycle*. 2018;17(1):53-63. doi:10.1080/15384101.2017.1387701
33. Estruel-Amades, S.; Massot-Cladera, M.; Pérez-Cano, F.J.; Franch, À.; Castell, M.; Camps-Bossacoma, M. Hesperidin Effects on Gut Microbiota and Gut-Associated Lymphoid Tissue in Healthy Rats. *Nutrients*. 2019;11(2):324. doi:10.3390/nu11020324
34. Havlik, J.; Marinello, V.; Gardyne, A.; et al. Dietary Fibres Differentially Impact on the Production of Phenolic Acids from Rutin in an In Vitro Fermentation Model of the Human Gut Microbiota. *Nutrients*. 2020;12(6):1577. doi:10.3390/nu12061577
35. Birt, D.F.; Hendrich, S.; Wang, W. Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacol Ther*. 2001;90(2-3):157-177. doi:10.1016/s0163-7258(01)00137-1
36. Doron, S.; Snyderman, D.R.; Gorbach, S.L. Lactobacillus GG: bacteriology and clinical applications. *Gastroenterol Clin North Am*. 2005;34(3):483-ix. doi:10.1016/j.gtc.2005.05.011
37. Segers, M.E.; Lebeer, S. Towards a better understanding of Lactobacillus rhamnosus GG-host interactions. *Microb Cell Fact*. 2014;13 Suppl 1(Suppl 1): S7. doi:10.1186/1475-2859-13-S1-S7
38. Foley, M.H.; O'Flaherty, S.; Allen, G.; et al. Lactobacillus bile salt hydrolase substrate specificity governs bacterial fitness and host colonization. *Proc Natl Acad Sci U S A*. 2021;118(6):e2017709118. doi:10.1073/pnas.2017709118