

 <https://doi.org/10.56238/alookdevelopv1-147>

Isabela Natália da Silva Ferreira

Graduate Program in Biotechnology. Northeast Biotechnology Network (RENORBIO) Federal Rural University of Pernambuco – UFRPE, 52171-900, Recife, Pe – Brazil
E-mail: isabelanatalia13@hotmail.com
ORCID: <https://orcid.org/0000-0002-2680-1143>

Galba Maria Campos-Takaki

Full Professor of the ICAM-TECH School. University Católica de Pernambuco – UNICAP, 50050-900 Recife-Pe, Brazil
E-mail: galba.takaki@unicap.br
ORCID: <https://orcid.org/0000-0002-0519-0849>

ABSTRACT

The literature describes several studies focused on the application of *S. maltophilia* in different areas of biotechnology. Due to its ease of interaction with numerous environments and living beings, this bacterium has an excellent adaptability, being able to convert different substrates into products of high environmental and technological interest.

Stenotrophomonas maltophilia is a gram-negative, aerobic, ubiquitous, rod-shaped bacterium belonging to the class of gammaproteobacteria and can be found in various natural and anthropogenic environments, e.g. soil, water, sediments or in association with plants. Biosurfactants are an alternative, compatible with the environment, for the use of chemically synthesized surfactants, since it is a bioproduct obtained through the metabolism of microorganisms and plants. Like their synthetic counterparts, biosurfactants comprise a wide variety of chemical structures and also excellent surfactant properties. The literature presents a wide range of studies focused on the production of biosurfactant by *S. maltophilia* using several substrates, temperature conditions, pH, salinade and fermentation times. In addition, there is also a huge variety of application of these biosurfactants produced, which drives the incentive to study this microorganism in the production of biosurfactants.

Keywords: Biodegradation, Biosurfactant, Environment, Bacteria.

1 INTRODUCTION

Surfactants are chemical molecules with diverse functions and properties, including surface tension reduction, emulsification, oil displacement, and micelle formation (Drakontis; Amin, 2020). They can be of synthetic or biological origin. Synthetic types are derived from petrochemistry and classified into five groups: anionic, non-ionic, cationic, amphoteric, and polymeric (Mohanty et al., 2021). Synthetic surfactants are expensive and their toxic effect can persist in the environment. These negative impacts have increased the search for alternative surfactants of biological origin, "biosurfactants". Biosurfactants are non-toxic, biodegradable and compatible with the environment. The market value of commercially available biosurfactants is expected to reach \$2.6 billion by 2023, gaining more than 8% and generating 540 kilotons of supply by 2024 (Gaur et al., 2022b, Kumar et al., 2021). Biosurfactants are classified according to their biological origin, mode of action, and molecular weight and generally have unique properties that make them better alternatives to synthetic surfactants (Mgbechidinma et al., 2022).

An integrated green economy for the production of biosurfactants overcomes the challenges associated with dispersed waste and the entire production chain (Johann et al., 2022). An integrated production system replaces the "take, make and dispose" structure with the concept of "resource, recovery and recycling" using natural by-products. Green production delays the use of virgin resources and reduces the extra production of waste from various sectors (Awasthi et al., 2022).

Industrial agro residues rich in fat and oil are raw materials of bioprocesses suitable for the production of biosurfactants, examples are the hydrolysate of animal fat and the residues of the palm oil refinery. There are valuable substrates for a multitude of fermenting microbes, including cooking oil (Shi et al., 2021), rice bran oil residues (Mishra et al., 2021), and plant-derived oils (sunflower, soybean, rapeseed, corn, and babassu oils) (Joy et al., 2019; Patowary et al., 2016; Radzuan et al., 2017). Oily residues offer low-cost lipid substrates with precise lipid-carbohydrate balance for microbial growth (Ibrahim et al., 2021; Mgbechidinma et al., 2023; Sohail; Jamil, 2020). Generally, the nutrient content in food waste is heterogeneous, containing about 40 to 60% starch; 5–10% protein; 0.84–1.38% salt; and 10-40% fatty or oily content (Demichelis et al., 2017). Due to their large volumes, low costs and sustainable sources of carbon and nitrogen for microbial growth, agro-industrial residues are considered green substrates for the production of biosurfactants. In addition to assisting in the management of environmental waste, the use of agri-food waste as a substrate also reduces the costs of producing biosurfactants (Das; Kumar, 2018; Mohanty et al., 2021; Sondh et al., 2022).

In this sense, the present study addresses several types of research focused on the production of biosurfactants obtained from the bacterium *Stenotrophomonas maltophilia*. In addition to presenting studies focused on the application of these biomolecules of high biotechnological, industrial and environmental interest.

2 STENOTROPHOMONAS MALTOPHILIA

Stenotrophomonas maltophilia is a gram-negative, aerobic, ubiquitous, rod-shaped bacterium belonging to the class of gammaproteobacteria (Chauviat; Meyer; Favre-Bonté, 2023). It is found in various natural and anthropogenic environments, e.g. soil, water, sediments or in association with plants (Ryan et al., 2009). *S. maltophilia* is also found in clinical settings where it can cause a wide variety of infections (Denton et al., 1998). The main studies about this bacterium are related to the clinical area. This opportunistic pathogen has low virulence but is of particular interest due to its natural and intrinsic phenotype of antibiotic resistance (Sanz-García et al., 2021; Deredjian et al., 2016).

S. maltophilia plays important ecosystem functions in sulfur and nitrogen cycles, in the degradation of complex compounds and pollutants, and in promoting plant growth and health. *Stenotrophomonas* can also colonize extreme artificial niches in hospitals, space shuttles, and clean

rooms. *S. maltophilia* has emerged as a global opportunistic human pathogen, which generally does not infect healthy hosts, but is associated with high morbidity and mortality in severely immunocompromised and debilitated individuals (Na; Berg, 2018).

2.1 OCCURRENCE OF *S. MALTOPHILIA* IN THE ENVIRONMENT

2.1.1 Soils, water, aquatic environments, sediments

S. maltophilia is found in several soils around the world (Youenou et al., 2015; Abdelrazek et al., 2020; Furlan et al., 2019; Javaid et al., 2020; Shreya et al., 2020). Their presence has been established in soils polluted with various types of contaminants such as heavy metals and antibiotics (Youenou et al., 2015), heavy metals from a mine (Liaquat et al., 2020), metals and industrial minerals from a mineral mine (Arulazhagan et al., 2017). It has also been found in the soil of a gold mine (Nangia et al., 2009) and in soils polluted with zinc and lead from garbage piles (Oleńska et al., 2020), cadmium and chromium (Chien et al., 2007; Chen et al., 2022) and copper (Andreazza et al., 2012). It has been recovered in soils contaminated with crude oil, transformer oil, and sulfurous oil (Elufisan et al., 2019; 2020; Somaraja et al., 2013), soils contaminated with fuel, diesel, gasoline, hydrocarbons and oil (Shahzad et al., 2016) and chemical plants contaminated soil (Lara-Moreno et al., 2021). It has been found in soils containing industrial wine residues (Snellinx et al., 2003), pesticides and surfactants of alkylphenol polyethoxylate (Kumar et al., 2007), an area of e-waste dismantling (Chen et al., 2013) or even plastic debris (Frederico et al., 2021).

The bacterium *S. maltophilia* has been found in soils where the presence of a contaminant is not specified, including bulk soil (Juhnke et al., 1989), unforested dunes of the Merzouga desert (Chemao-Elfihri et al., 2020), drought-prone ecosystems (Haque et al., 2020), beach sand (Singh et al., 2011), soil of an island (Park et al., 2013), soil on the banks of a river (Dubey et al., 2013), soil near a glacier (Kuddus; Ramteke, 2009) muddy soil (Miyaji et al., 2005) and slaughterhouse soil (Waghmare et al., 2015). It is present in soils associated with the use of agricultural land, such as cultivated and altered soils, but also in alterations (Patil et al., 2016; Chauviat; Meyer; Favre-Bonté, 2023).

S. maltophilia has been identified in water, aquatic environments and sediments, its presence has been demonstrated in several natural aquatic environments such as rivers (Li, et al., 2019), sediments and water from rivers and streams (Ochoa-Sánchez; Vinuesa, 2017) on the surface of a lake (De La Rosa-García et al., 2007), in droplets of water from a cave (Enyedi et al., 2020), swamp water (Singh et al., 2011), seawater (Dewi et al., 2020), polluted coastal water (Mançano et al., 2020), groundwater (Martin et al., 2018), and biofilms in natural and artificial aquifers (Jägevall; Rabe; Pedersen, 2011).

In addition, *S. maltophilia* has been isolated in association with wastewater and water treatment plants, in surface water, groundwater, and bioremediation reactor samples from the Niagara watershed (Nakatsu et al., 1995), water treatment and subsequent distribution systems (Hoefel et al., 2005), a drinking water treatment plant (Gomes et al., 2020), wastewater treatment plants (Permala et al., 2018), effluents from palm oil mills (Bala et al., 2018), final effluents from a pig farm (Kim; Park; Seo, 2018) wastewater from an oil reservoir (Hassanshahian et al., 2013) and in contaminated sewage (Elufisan et al., 2020). Several types of sediments containing *S. maltophilia* have been identified, i.e., sewage sediments from agricultural drainage ponds (Dungan et al., 2003), anaerobic and activated sludge from a wastewater treatment plant (Chen et al., 2020; URSZULA et al., 2009; Xiong et al., 2020), wastewater bioreactor sludge (Bramucci; Nagarajan, 2000) wastewater sediments in a processing unit (Al-Dhabi et al., 2021), sediment from a coal tailings solids storage facility (Siddique et al., 2007), coastal sediment from a shipyard (Patel; Patel; Datta, 2013), sediment of a channel (DUDÁŠOVÁ et al., 2014), fresh water (Jayaraman; Sigamani; Ramamurthy, 2021), seawater (Jamal et al., 2018), and polluted river (Iyer; Damania; Iken, 2019), sediments, and even microplastics from a river (ZHANG, et al. 2021).

2.1.2 Interaction with plants and animals

S. maltophilia interacts with several plants, notably in the rhizosphere. It has been isolated from the rhizospheres of landfill-growing plants (Franco et al., 2005), rice (*Oryza sativa*) (Mir et al., 2022), oats (*Avena sativa*) and plantains (*Plantago lanceolata*) (Andreazza et al., 2022), cabbage (*Brassica oleracea*), mustard (*Sinapis alba*), corn (*Triticum vulgare*), beetroot (*Beta vulgaris*) (Dunne et al., 1997), potato (*Solanum tuberosum*) (Lottmann et al., 1999), tomato plants (*Solanum lycopersicum*) (Haque et al., 2020), eggplant (*Solanum melongena*) (Messiha et al., 2007) and strawberry (*Fragaria x ananassa*) (Berg et al., 2002). Strains were recovered from moist brown forest soil (Krsek; Wellington, 2001) and grassland. *S. maltophilia* is part of the phyllosphere of green beans (*Phaseolus vulgaris*) (Leng et al., 2016), potato (*Solanum tuberosum*) (Wilson; Lindow, 1994) and rice (*Oryza sativa*) and has been described as an epiphyte of common ivy (*Hedera helix*) (Sahu et al., 2021), as well as several other plant and plant species.

S. maltophilia is found in reptiles, for example, in the oral microflora of snakes (Hejnar et al., 2007) and in prairie rattlesnakes (Fitzgerald et al., 2013). It has been blamed for chronic respiratory disease of the king python (Albini et al., 2009) and septicemia in the dwarf crocodile of West Africa (Harris; Rogers, 2001). *S. maltophilia* was isolated from mammals. It has been found in the feces of tapirs (Guan et al., 2008), in porcine semen (Althouse et al., 2000) and can cause mastitis in cows (Bhandari et al., 2014), chronic respiratory disease in horses, cats and dogs. *S. maltophilia* has been

detected in association with fish, including cultured yellowtail (*Seriola quinqueradiata*) (FURUSHITA et al., 2005), cultivated channel catfish (*Ictalurus punctatus*) (Geng et al., 2010), in the kidneys of juvenile African catfish (*Clarias gariepinus*) (Abraham et al., 2016). and in the spleen of rainbow trout (*Oncorhynchus mykiss*) (Duman; Saticioglu; Altun, 2019) where it causes disease. In birds, *S. maltophilia* has been recovered from feces on a poultry farm (Hassan et al., 2021), from the environment of poultry flocks (Shah et al., 2020), and from the gut microbiota of Condor (Jacobs et al., 2019).

3 BIOSSURFACTANTS

Environmental legislation has evolved rapidly to develop sustainable products. This has inspired modern researchers to examine natural resources for industrial and domestic applications (Othmani et al., 2022; Akcil et al., 2015). Among several synthetic products, surfactants or surfactants are widely used worldwide, with a growing trend of their synthesis and subsequent application (Inès; Dhouha, 2015). Due to their distinct properties, e.g., dispersion, emulsification, biological activity, surfactants have found several industrial applications (Shahid et al., 2021; Banat et al., 2021). Chemical surfactants, also known as surfactants, are synthetic molecules used primarily in industrial products such as washing powder, detergents, toothpastes, soaps, and shampoos (Onaizi, 2021; Zou et al., 2014; Patel; Patel, 2020.).

Surfactants are chemical compounds formed by amphipathic molecules containing hydrophilic and hydrophobic portions that divide into physical interfaces. Apolar portions are often hydrocarbon chains, while polar portions can be cationic, anionic, non-ionic, or amphoteric molecules (Sarubbo et al., 2022). This combination of hydrophobic and hydrophilic portions allows surfactants to reduce surface and interfacial stresses and form microemulsions, in which hydrocarbons are solubilized in water or vice versa (Farias et al., 2021). The most effective way to characterize a surfactant is by measuring the force of attraction between liquid molecules, thus classifying the surfactant in its ability to affect surface and interfacial tensions. Effective surfactants reduce surface tensions, facilitating interactions between molecules of different polar natures (Santos et al., 2016). Most commercially available surfactants are derived from petroleum-based organic compounds, so scientists are now eager for their possible replacement. However, its implication is associated with various environmental concerns due to its source of origin (Kashif et al., 2022). Some recent scientific evidence has suggested the use of bio-based surface active compounds as an innovative and green replacement of their counterparts (Ambaye et al., 2021; Hentati et al., 2021).

The biosurfactant is an alternative, compatible with the environment, for the use of chemically synthesized surfactants since it is a bioproduct obtained through the metabolism of microorganisms

and plants. Like their synthetic counterparts, biosurfactants comprise a wide variety of chemical structures. They are produced by microorganisms grown on insoluble (oils, residues and hydrocarbons) or soluble (carbohydrates) substrates (Silva et al., 2014; Bezerra et al., 18). Replacing synthetic surfactants with biosurfactants would reduce CO₂ emissions by 8%, avoiding about 1.5 million tons of CO₂ released into the atmosphere (Farias et al., 2021; Rocha and Silva et al., 2021; Banat et al., 2021). Currently, biosurfactants occupy about 10% of the total world production of surfactants (approximately ten million tons per year) (Sarubbo et al., 2022). These natural surfactants have applications in the petroleum, food (as emulsifiers), pharmaceutical (formulation of moisturizers, creams and medicines), medical (antimicrobial agents), agricultural (fertilizers) and civil (waste and sewage treatment) industries (Ribeiro; War; Sarubbo, 2020).

4 APPLICATIONS OF *STENOTROPHOMONAS MALTOPHILIA*

4.1 BIOSURFACTANT PRODUCTION

The literature presents a wide range of studies focused on the production of biosurfactant by *S. maltophilia* using several substrates, temperature conditions, pH, salinade and fermentation times. In addition, there is also a huge variety of application of these biosurfactants produced, which drives the incentive to study this microorganism in the production of biosurfactants.

In a study conducted by Ferreira et al., (2022) the bacterium *S. maltophilia* UCP 1601 converted the low-cost renewable substrates, milhocin, residual glycerol and post-frying soybean oil into an anionic biosurfactant with surface tension of 27mN/m, in addition to stability in different environmental conditions, such as temperature, salinity and pH. All the determinations made confirm the generation of a stable and efficient bioproduct that contributes to the circular bioeconomy and the great potential for remediation in adverse environmental conditions.

Larik et al (2019) evaluated biosurfactant production by *S. maltophilia*. The bacterium showed excellent potential for the production of biosurfactant (oil displacement assay=>6 mm; surface tension=33.9 and 34.4 m N m⁻¹) and lipase enzyme (lipolytic zone=1.8±0.1 cm and 2.2±0.3 cm).

The study conducted by Ezebuoro et al (2019) investigated the effects of nitrogen and carbon sources on the production of biosurfactant by a bacterium that uses hydrocarbons as a substrate by the bacterium, *Stenotrophomonas* sp. The surface tension obtained was 30.85 mN/m. The authors highlight that this method of biosurfactant production can be extended for industrial purposes.

4.2 BIODEGRADATION AND BIOSURFACTANT PRODUCTION

The process of biodegradation happens when the structure of contaminating compounds is transformed with the breakdown of their molecules, in a biotic way, making them, in most cases, less toxic than the original molecules (Fenner et al., 2013; Oliveira et al., 2022).

A study conducted by Larik et al (2019) evaluated the biodegradation of two petrochemical hydrocarbons, namely diesel oil and motor oil used by *S. maltophilia* with blackchemical substrates as the only carbon source. The biodegradation rates of hydrocarbons were estimated in terms of % removal using UV-Vis spectrophotometer, while the characteristic spectral changes in both hydrocarbons were confirmed using infrared Fourier transform spectrometry with attenuated total reflection. Significant degradation rates of 88.5 and 71.8% (respectively) were achieved during the experiment.

Aliyu et al (2022) investigated the potential use of bacterial isolates expressing biosurfactants in the biodegradation of one of the products of the petroleum refining process (PMS- Premium Motor Spirit). Among the microorganisms tested, *S. maltophilia* showed excellent biosurfactant production results in the biodegradation process. The authors point out that the ability of these isolates to grow in mineral salt media supplemented with PMS as the only carbon source presents a true path for exploration in biotechnology, towards bioremediation mediated by biosurfactants of hydrocarbon pollutants in oil-contaminated soils.

A study by Patel, Patel (2020) evaluated the biosurfactant production capacity of *Stenotrophomonas* sp. nov. S1VKR-26, profiling its bioremediation capacity to remediate wastewater from oil refineries in a laboratory-scale bioreactor and evaluating of the phytotoxicity of bioremediated petroleum wastewater. Different PAHs (Polycyclic Aromatic Hydrocarbons) such as naphthalene (93%), phenanthrene (86%), fluoranthene (92%) and pyrene (98.3%), total petroleum hydrocarbons (72.33%) and phenolic compounds (93.06%) were significantly remediated from wastewater after treatment of the S1VKR-26 strain. In addition, the diluted petroleum effluent 1:1 treated with the S1VKR-26 strain has higher germination (100%), vigor (486) and seedling (4.86 cm) compared to the untreated effluent. Indicating an excellent potential of biosurfactant production in the biodegradation process.

Studies indicate the genus *Stenotrophomonas* as a producer of biosurfactant in the process of engine oil degradation (Gargouri et al., 201; Larik et al., 2019), crude oil degradation (Muthukamalam et al., 2017), and diesel oil degradation (Larik et al., 2019). A study by Deng et al. (2015) points to the degradation of eight organophosphate pesticides by a strain of *Stenotrophomonas maltophilia* in 24 h (Deng et al., 2015). Nogueira et al (2020) conducted a study with the bacterium *Stenotrophomonas maltophilia* in the production of bioemulsifier, having post-frying soybean oil as a source of insoluble

carbon, this biomolecule was characterized of polymeric nature and obtained excellent Emulsification Index with burnt motor oil of 82.74%. The biosurfactant of *S. maltophilia* was reported as an iron chelating agent in a study by Hemlata, Selvin and Tukaram (2015).

A study by Zaffar et al, (2018) described the degradation of the organochlorine pesticide endosulfan by a strain of *Stenotrophomonas maltophilia* isolated from contaminated soil. The bacterium demonstrated the ability to use endosulfan as the sole source of sulfur and excellent pesticide degradation capacity.

4.3 ENZYME PRODUCTION

The literature also describes the production of enzymes by this bacterium (Muthukamalam et al., 2017; Wang et al., 2016). Aktas et al (2022) evaluated the production of chitinase by *S. maltophilia* for application as a biological control agent. The enzyme produced was effective in reducing the growth of the fungus *Fusarium oxysporum* and in destroying the chitin structure of the potato beetle.

The study by Subramanian et al (2022) summarizes the isolation of chitinase-producing *S. maltophilia* bacteria from different shrimp waste disposal sites in Parangipettai (India). After purification, the specific activity of chitinase was 5.01 (U/ml) and the protein content was 72 mg and the recovery rate was 48.06%. The results of the study cover that the use of foods enriched with chitin derivatives is taken to large-scale approaches, thus benefiting the aquaculture sector.

The bacterium capable of producing extracellular fibrinolytic protease was isolated and identified as *Stenotrophomonas maltophilia* Gd2 based on ribotyping. The in vitro fibrinolytic profile of this enzyme showed 73% fibrin clot dissolution in 4h. Further optimization of enzyme production was achieved with the implementation of the Plackett-Burman experimental design, where production levels increased to 3411FUml-1 and noted that peptone, pH, dextrose and K₂HPO₄ were considered significant factors. The author describes that this result reports the higher yield of fibrinolytic enzyme with *S. maltophilia* about reports in the literature.

REFERENCES

- ABDELRAZEK, Nada A. et al. Production, characterization and bioinformatics analysis of L-asparaginase from a new *Stenotrophomonas maltophilia* EMCC2297 soil isolate. *AMB Express*, v. 10, n. 1, p. 71, 2020.
- ABRAHAM, T. Jawahar et al. *Stenotrophomonas maltophilia* as an opportunistic pathogen in cultured African catfish *Clarias gariepinus* (Burchell, 1822). *Aquaculture*, v. 450, p. 168-172, 2016.
- AKCIL, Ata et al. A review of approaches and techniques used in aquatic contaminated sediments: metal removal and stabilization by chemical and biotechnological processes. *Journal of Cleaner Production*, v. 86, p. 24-36, 2015.
- AKTAS, Cigdem et al. Purification and characterization of *Stenotrophomonas maltophilia* chitinase with antifungal and insecticidal properties. *Preparative Biochemistry & Biotechnology*, p. 1-10, 2022.
- ALBINI et al. *Stenotrophomonas maltophilia* isolated from the airways of animals with chronic respiratory disease. *Schweizer Archiv für Tierheilkunde*, v. 151, n. 7, p. 323-328, 2009.
- AL-DHABI, Naif Abdullah et al. Removal of nitrogen from wastewater of date processing industries using a Saudi Arabian mesophilic bacterium, *Stenotrophomonas maltophilia* Al-Dhabi-17 in sequencing batch reactor. *Chemosphere*, v. 268, p. 128636, 2021.
- ALIYU, Aminu et al. Biodegradation of Premium Motor Spirit using Surfactant-expressing Bacteria from Mechanic Workshops in Malumfashi, Katsina State, Nigeria.
- ALTHOUSE, G. C. et al. Field investigations of bacterial contaminants and their effects on extended porcine semen. *Theriogenology*, v. 53, n. 5, p. 1167-1176, 2000.
- AMBAYE, Teklit Gebregiorgis et al. Preparation, characterization and application of biosurfactant in various industries: A critical review on progress, challenges and perspectives. *Environmental Technology & Innovation*, v. 24, p. 102090, 2021.
- AN, Shi-qi; BERG, Gabriele. *Stenotrophomonas maltophilia*. *Trends in microbiology*, v. 26, n. 7, p. 637-638, 2018.
- ANDREAZZA, Robson et al. Characterization of copper-resistant rhizosphere bacteria from *Avena sativa* and *Plantago lanceolata* for copper bioreduction and biosorption. *Biological Trace Element Research*, v. 146, p. 107-115, 2012.
- ARULAZHAGAN, Pugazhendi et al. Biodegradation of polycyclic aromatic hydrocarbons by an acidophilic *Stenotrophomonas maltophilia* strain AJH1 isolated from a mineral mining site in Saudi Arabia. *Extremophiles*, v. 21, p. 163-174, 2017.
- AWASTHI, Mukesh Kumar et al. Agricultural waste biorefinery development towards circular bioeconomy. *Renewable and Sustainable Energy Reviews*, v. 158, p. 112122, 2022.
- BALA, Jeremiah David et al. Microbiota of palm oil mill wastewater in Malaysia. *Tropical life sciences research*, v. 29, n. 2, p. 131, 2018.

BANAT, Ibrahim M. et al. Biosurfactants: The green generation of speciality chemicals and potential production using Solid-State fermentation (SSF) technology. *Bioresource Technology*, v. 320, p. 124222, 2021.

BERG, Gabriele et al. Plant-dependent genotypic and phenotypic diversity of antagonistic rhizobacteria isolated from different *Verticillium* host plants. *Applied and environmental microbiology*, v. 68, n. 7, p. 3328-3338, 2002.

BEZERRA, Káren Gercyane O. et al. Saponins and microbial biosurfactants: Potential raw materials for the formulation of cosmetics. *Biotechnology progress*, v. 34, n. 6, p. 1482-1493, 2018.

BHANDERI, Bharat B. et al. Cultural and metagenomic based identification of a microbiome from subclinical mastitis in cows. *Veterinarski arhiv*, v. 84, n. 3, p. 215-228, 2014.

BRAMUCCI, Michael G.; NAGARAJAN, Vasantha. Industrial wastewater bioreactors: sources of novel microorganisms for biotechnology. *Trends in biotechnology*, v. 18, n. 12, p. 501-505, 2000.

CHAUVIAT, Amandine; MEYER, Thibault; FAVRE-BONTÉ, Sabine. Versatility of *Stenotrophomonas maltophilia*: Ecological roles of RND efflux pumps. *Heliyon*, 2023.

CHEMAO-ELFIHRI, M. W. et al. Draft genome sequence of *Stenotrophomonas maltophilia* MDMC339, isolated from soil of Merzouga desert in Morocco. *Microbiology Resource Announcements*, v. 9, n. 32, p. e00634-20, 2020.

CHEN, Shuona et al. Effect of copper (II) on biodegradation of benzo [a] pyrene by *Stenotrophomonas maltophilia*. *Chemosphere*, v. 90, n. 6, p. 1811-1820, 2013.

CHEN, Wenfang et al. Isolation of functional bacterial strains from chromium-contaminated site and bioremediation potentials. *Journal of Environmental Management*, v. 307, p. 114557, 2022.

CHIEN, Chih-Ching; HUNG, Chun-Wei; HAN, Chu-Ting. Removal of cadmium ions during stationary growth phase by an extremely cadmium-resistant strain of *Stenotrophomonas* sp. *Environmental Toxicology and Chemistry: An International Journal*, v. 26, n. 4, p. 664-668, 2007.

DAS, Amar Jyoti; KUMAR, Rajesh. Utilization of agro-industrial waste for biosurfactant production under submerged fermentation and its application in oil recovery from sand matrix. *Bioresource technology*, v. 260, p. 233-240, 2018.

DE LA ROSA-GARCÍA, S. C. et al. Antimicrobial properties of moderately halotolerant bacteria from cenotes of the Yucatan peninsula. *Letters in applied microbiology*, v. 45, n. 3, p. 289-294, 2007.

DEMICHELIS, Francesca et al. Investigation of food waste valorization through sequential lactic acid fermentative production and anaerobic digestion of fermentation residues. *Bioresource technology*, v. 241, p. 508-516, 2017.

DENG, Shuyan et al. Rapid biodegradation of organophosphorus pesticides by *Stenotrophomonas* sp. G1. *Journal of hazardous materials*, v. 297, p. 17-24, 2015.

DENTON, Miles; KERR, Kevin G. Microbiological and clinical aspects of infection associated with *Stenotrophomonas maltophilia*. *Clinical microbiology reviews*, v. 11, n. 1, p. 57-80, 1998.

DEREDJIAN, Amélie et al. Occurrence of *Stenotrophomonas maltophilia* in agricultural soils and antibiotic resistance properties. *Research in microbiology*, v. 167, n. 4, p. 313-324, 2016.

DEWI, Dewa AP Rasmika; GÖTZ, Barbara; THOMAS, Torsten. Diversity and genetic basis for carbapenem resistance in a coastal marine environment. *Applied and Environmental Microbiology*, v. 86, n. 10, p. e02939-19, 2020.

DUBEY, Kriti Kumari; FULEKAR, M. H. Investigation of potential rhizospheric isolate for cypermethrin degradation. *3 Biotech*, v. 3, p. 33-43, 2013.

DUDÁŠOVÁ, Hana et al. Bacterial strains isolated from PCB-contaminated sediments and their use for bioaugmentation strategy in microcosms. *Journal of Basic Microbiology*, v. 54, n. 4, p. 253-260, 2014.

DUMAN, Muhammed; SATICIOGLU, Izzet Burcin; ALTUN, Soner. Molecular characterization and antimicrobial resistance profile of fecal contaminants and spoilage bacteria that emerge in rainbow trout (*Oncorhynchus mykiss*) farms. *Bioscience of Microbiota, Food and Health*, v. 38, n. 2, p. 41-48, 2019.

DUNNE, Colum et al. Biological control of *Pythium ultimum* by *Stenotrophomonas maltophilia* W81 is mediated by an extracellular proteolytic activity. *Microbiology*, v. 143, n. 12, p. 3921-3931, 1997.

DUNGAN, Robert S.; YATES, Scott R.; FRANKENBERGER JR, William T. Transformations of selenate and selenite by *Stenotrophomonas maltophilia* isolated from a seleniferous agricultural drainage pond sediment. *Environmental Microbiology*, v. 5, n. 4, p. 287-295, 2003.

ELUFISAN, Temidayo O. et al. Antimicrobial susceptibility pattern of *Stenotrophomonas* species isolated from Mexico. *African Health Sciences*, v. 20, n. 1, p. 168-181, 2020.

ELUFISAN, Temidayo Oluyomi et al. Complete genome sequence of *Stenotrophomonas maltophilia* strain SVIA2, isolated from crude oil-contaminated soil in Tabasco, Mexico. *Microbiology Resource Announcements*, v. 8, n. 30, p. e00529-19, 2019.

ENYEDI, Nóra Tünde et al. Cave bacteria-induced amorphous calcium carbonate formation. *Scientific reports*, v. 10, n. 1, p. 8696, 2020.

EZEBUIRO, Victor et al. Effects of Nitrogen and Carbon Sources on Biosurfactant Production by Hydrocarbon-utilizing *Stenotrophomonas* sp. *Microbiology Research Journal International*, v. 29, n. 5, p. 1-10, 2019.

FARIAS, Charles Bronzo B. et al. Production of green surfactants: Market prospects. *Electronic Journal of Biotechnology*, v. 51, p. 28-39, 2021.

FENNER, Kathrin et al. Evaluating pesticide degradation in the environment: blind spots and emerging opportunities. *science*, v. 341, n. 6147, p. 752-758, 2013.

FERREIRA, Isabela Natália da Silva et al. Production and functional stability of the biourfactant isolated from *Stenotrophomonas maltophilia* UCP 1601. *Research, Society and Development*, v. 11, n. 11, p. e87111132995-e87111132995, 2022.

FITZGERALD, Kevin T. et al. Additional observations and notes on the natural history of the prairie rattlesnake (*Crotalus viridis*) in Colorado. *Topics in companion animal medicine*, v. 28, n. 4, p. 167-176, 2013.

FRANCO, A. R. et al. Isolation and characterization of polymeric galloyl-ester-degrading bacteria from a tannery discharge place. *Microbial ecology*, v. 50, p. 550-556, 2005.

FREDERICO, Tayná Diniz et al. Draft Genome Sequence of *Stenotrophomonas maltophilia* Strain PE591, a Polyethylene-Degrading Bacterium Isolated from Savanna Soil. *Microbiology resource announcements*, v. 10, n. 32, p. e00490-21, 2021.

FURLAN, João Pedro Rueda et al. Characterization of acquired antimicrobial resistance genes in environmental *Stenotrophomonas maltophilia* isolates from Brazil. *Microbial Drug Resistance*, v. 25, n. 4, p. 475-479, 2019.

FURUSHITA, Manabu et al. Isolation of multidrug-resistant *Stenotrophomonas maltophilia* from cultured yellowtail (*Seriola quinqueradiata*) from a marine fish farm. *Applied and environmental microbiology*, v. 71, n. 9, p. 5598-5600, 2005.

GARGOURI, Boutheina et al. Biosurfactant production by the crude oil degrading *Stenotrophomonas* sp. B-2: chemical characterization, biological activities and environmental applications. *Environmental Science and Pollution Research*, v. 24, p. 3769-3779, 2017.

GENG, Yi et al. *Stenotrophomonas maltophilia*, an emerging opportunist pathogen for cultured channel catfish, *Ictalurus punctatus*, in China. *Aquaculture*, v. 308, n. 3-4, p. 132-135, 2010.

GOMES, Thiago Santos et al. Presence and interaction of free-living amoebae and amoeba-resisting bacteria in water from drinking water treatment plants. *Science of the Total Environment*, v. 719, p. 137080, 2020.

GUAN, Shu et al. Aflatoxin B1 degradation by *Stenotrophomonas maltophilia* and other microbes selected using coumarin medium. *International journal of molecular sciences*, v. 9, n. 8, p. 1489-1503, 2008.

HAQUE, Md Manjurul et al. Biofilm producing rhizobacteria with multiple plant growth-promoting traits promote growth of tomato under water-deficit stress. *Frontiers in Microbiology*, v. 11, p. 542053, 2020.

HARRIS, N. Beth; ROGERS, Douglas G. Septicemia associated with *Stenotrophomonas maltophilia* in a West African dwarf crocodile (*Osteolaemus tetraspis* subsp. *tetraspis*). *Journal of veterinary diagnostic investigation*, v. 13, n. 3, p. 255-258, 2001.

HASSANSHAHIAN, Mehdi et al. Isolation and characterization of alkane degrading bacteria from petroleum reservoir waste water in Iran (Kerman and Tehran provenances). *Marine pollution bulletin*, v. 73, n. 1, p. 300-305, 2013.

HEJNAR, Petr et al. *Stenotrophomonas maltophilia* as a part of normal oral bacterial flora in captive snakes and its susceptibility to antibiotics. *Veterinary microbiology*, v. 121, n. 3-4, p. 357-362, 2007.

Hemlata, B., Selvin, J., & Tukaram, K. (2015). Optimization of iron chelating biosurfactant production by *Stenotrophomonas maltophilia* NBS-11. *Biocatalysis and agricultural biotechnology*, 4(2), 135-143.

HENTATI, Dorra et al. Investigation of halotolerant marine *Staphylococcus* sp. CO100, as a promising hydrocarbon-degrading and biosurfactant-producing bacterium, under saline conditions. *Journal of Environmental Management*, v. 277, p. 111480, 2021.

HOEFEL, Daniel et al. Profiling bacterial survival through a water treatment process and subsequent distribution system. *Journal of applied microbiology*, v. 99, n. 1, p. 175-186, 2005.

IBRAHIM, Sami; DIAB, Atef; ABDULLA, Hesham. Bio-cleaning Efficiency of Rhamnolipids Produced from Native *Pseudomonas aeruginosa* Grown on Agro-industrial By-products for Liquid Detergent Formulation. *Applied Biochemistry and Biotechnology*, v. 193, p. 2616-2633, 2021.

INÈS, Mnif; DHOUHA, Ghribi. Glycolipid biosurfactants: Potential related biomedical and biotechnological applications. *Carbohydrate Research*, v. 416, p. 59-69, 2015.

IYER, Rupa; DAMANIA, Ashish; IKEN, Brian. Genome data of *Stenotrophomonas maltophilia* DF07 collected from polluted river sediment reveals an opportunistic pathogen and a potential antibiotic reservoir. *Data in brief*, v. 25, 2019.

JÄGEVALL, Sara; RABE, Lisa; PEDERSEN, Karsten. Abundance and diversity of biofilms in natural and artificial aquifers of the Äspö Hard Rock Laboratory, Sweden. *Microbial ecology*, v. 61, p. 410-422, 2011.

JAVAID, Hadiqa et al. Biosynthesis of polyhydroxyalkanoates (PHAs) by the valorization of biomass and synthetic waste. *Molecules*, v. 25, n. 23, p. 5539, 2020.

JACOBS, Lindsey et al. California condor microbiomes: Bacterial variety and functional properties in captive-bred individuals. *PLoS One*, v. 14, n. 12, p. e0225858, 2019.

JAYARAMAN, Jamunadevi; SIGAMANI, Santhosh; RAMAMURTHY, Dhandapani. Metal biosorption by magnetotactic bacteria isolated from fresh water sediments and characterization of extracted magnetosomes. *Archives of Microbiology*, v. 203, n. 10, p. 5951-5962, 2021.

JOHANN, Sarah et al. A plea for the integration of Green Toxicology in sustainable bioeconomy strategies—Biosurfactants and microgel-based pesticide release systems as examples. *Journal of Hazardous Materials*, v. 426, p. 127800, 2022.

JOY, Sam et al. Statistical and sequential (fill-and-draw) approach to enhance rhamnolipid production using industrial lignocellulosic hydrolysate C6 stream from *Achromobacter* sp.(PS1). *Bioresource technology*, v. 288, p. 121494, 2019.

JUHNKE, M. ELLIOTT; DES JARDIN, ELIZABETH. Selective medium for isolation of *Xanthomonas maltophilia* from soil and rhizosphere environments. *Applied and Environmental Microbiology*, v. 55, n. 3, p. 747-750, 1989.

KASHIF, Ayesha et al. Current advances in the classification, production, properties and applications of microbial biosurfactants—A critical review. *Advances in Colloid and Interface Science*, p. 102718, 2022.

KIM, Young-Ji; PARK, Jin-Hyeong; SEO, Kun-Ho. Presence of *Stenotrophomonas maltophilia* exhibiting high genetic similarity to clinical isolates in final effluents of pig farm wastewater treatment plants. *International Journal of Hygiene and Environmental Health*, v. 221, n. 2, p. 300-307, 2018.

KRSEK, Martin; WELLINGTON, Elizabeth MH. Assessment of chitin decomposer diversity within an upland grassland. *Antonie Van Leeuwenhoek*, v. 79, p. 261-267, 2001.

KUDDUS, Mohammed; RAMTEKE, Pramod W. Cold-active extracellular alkaline protease from an alkaliphilic *Stenotrophomonas maltophilia*: production of enzyme and its industrial applications. *Canadian journal of microbiology*, v. 55, n. 11, p. 1294-1301, 2009.

KUMAR, Koel et al. Enrichment and isolation of endosulfan degrading and detoxifying bacteria. *Chemosphere*, v. 68, n. 2, p. 317-322, 2007.

LARA-MORENO, Alba et al. A comprehensive feasibility study of effectiveness and environmental impact of PAH bioremediation using an indigenous microbial degrader consortium and a novel strain *Stenotrophomonas maltophilia* CPHE1 isolated from an industrial polluted soil. *Journal of Environmental Management*, v. 289, p. 112512, 2021.

LARIK, I. A. et al. *Stenotrophomonas maltophilia* strain 5DMD: an efficient biosurfactant-producing bacterium for biodegradation of diesel oil and used engine oil. *International Journal of Environmental Science and Technology*, v. 16, p. 259-268, 2019.

LENG, Yifei et al. Biotransformation of tetracycline by a novel bacterial strain *Stenotrophomonas maltophilia* DT1. *Journal of hazardous materials*, v. 318, p. 125-133, 2016.

LIAQUAT, Fiza et al. Cd-tolerant SY-2 strain of *Stenotrophomonas maltophilia*: a potential PGPR, isolated from the Nanjing mining area in China. *3 Biotech*, v. 10, p. 1-10, 2020.

LI, Dan et al. Isolation, characterization, and inactivation of *Stenotrophomonas maltophilia* from leafy green vegetables and urban agriculture systems. *Frontiers in Microbiology*, v. 10, p. 2718, 2019.

LOTTMANN, Jana et al. Influence of transgenic T4-lysozyme-producing potato plants on potentially beneficial plant-associated bacteria. *FEMS Microbiology Ecology*, v. 29, n. 4, p. 365-377, 1999.

MANÇANO, Stella Maria Casas Novas et al. Frequency and diversity of *Stenotrophomonas* spp. carrying blaKPC in recreational coastal waters. *Water research*, v. 185, p. 116210, 2020.

MARTIN, Misty S. et al. Characterization of bacterial diversity in contaminated groundwater using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Science of the Total Environment*, v. 622, p. 1562-1571, 2018.

MESSIHA, N. A. S. et al. *Stenotrophomonas maltophilia*: a new potential biocontrol agent of *Ralstonia solanacearum*, causal agent of potato brown rot. *European journal of plant pathology*, v. 118, p. 211-225, 2007.

MGBECHIDINMA, Chiamaka Linda et al. Fatty acid composition and nutritional analysis of waste crude fish oil obtained by optimized milder extraction methods. *Environmental Engineering Research*, v. 28, n. 2, 2023.

- MGBECHIDINMA, Chiamaka Linda et al. Integration of Green Economy Concepts for Sustainable Biosurfactant Production-A Review. *Bioresource Technology*, p. 128021, 2022.
- MIR, Mohammad Imran et al. Multifarious indigenous diazotrophic rhizobacteria of rice (*Oryza sativa* L.) rhizosphere and their effect on plant growth promotion. *Frontiers in nutrition*, v. 8, p. 781764, 2022.
- MISHRA, Ashutosh et al. Optimization of process inputs for the synthesis of waste rice bran oil isolated *Pseudomonas aeruginosa* MTCC 424 biosurfactant using response surface methodology for oil recovery applications. *Bioresource Technology Reports*, v. 14, p. 100653, 2021.
- MIYAJI, T. et al. Purification and characterization of extracellular alkaline serine protease from *Stenotrophomonas maltophilia* strain S-1. *Letters in applied microbiology*, v. 41, n. 3, p. 253-257, 2005.
- MORALES PEDRAZA, J. General Overview of the Energy Sector in the North America Region. *Conventional Energy in North America*, p. 1-87, 2019.
- MOHANTY, Swayansu Sabyasachi et al. A critical review on various feedstocks as sustainable substrates for biosurfactants production: a way towards cleaner production. *Microbial cell factories*, v. 20, n. 1, p. 1-13, 2021.
- MUTHUKAMALAM, Santhakumar et al. Characterization of dioxygenases and biosurfactants produced by crude oil degrading soil bacteria. *Brazilian journal of microbiology*, v. 48, p. 637-647, 2017.
- NANGIA, Yogesh et al. Facile biosynthesis of phosphate capped gold nanoparticles by a bacterial isolate *Stenotrophomonas maltophilia*. *Applied Physics Letters*, v. 94, n. 23, p. 233901, 2009.
- NAKATSU, C. H. et al. The phylogenetic distribution of a transposable dioxygenase from the Niagara River watershed. *Molecular ecology*, v. 4, n. 5, p. 593-604, 1995.
- NG, Yan Jer et al. Recent advances of biosurfactant for waste and pollution bioremediation: Substitutions of petroleum-based surfactants. *Environmental Research*, v. 212, p. 113126, 2022.
- NOGUEIRA, Ianne Batista et al. Bioconversion of agroindustrial waste in the production of bioemulsifier by *Stenotrophomonas maltophilia* UCP 1601 and application in bioremediation process. *International Journal of Chemical Engineering*, v. 2020, p. 1-9, 2020.
- OCHOA-SÁNCHEZ, Luz E.; VINUESA, Pablo. Evolutionary genetic analysis uncovers multiple species with distinct habitat preferences and antibiotic resistance phenotypes in the *Stenotrophomonas maltophilia* complex. *Frontiers in microbiology*, v. 8, p. 1548, 2017.
- OLEŃSKA, Ewa et al. *Trifolium repens*-associated bacteria as a potential tool to facilitate phytostabilization of zinc and lead polluted waste heaps. *Plants*, v. 9, n. 8, p. 1002, 2020.
- OLIVEIRA, Mayane et al. Avaliação da importância do uso de microrganismos na biodegradação de agrotóxico aplicados na agricultura—uma revisão. *Conjecturas*, v. 2022, n. 18, p. 368-379, 2022.
- ONAIZI, Sagheer A. Statistical analyses of the effect of rhamnolipid biosurfactant addition on the enzymatic removal of Bisphenol A from wastewater. *Biocatalysis and Agricultural Biotechnology*, v. 32, p. 101929, 2021.

OTHMANI, Amina et al. Agricultural waste materials for adsorptive removal of phenols, chromium (VI) and cadmium (II) from wastewater: A review. *Environmental Research*, v. 204, p. 111916, 2022.

PARK, Jong Myong; PARK, Sung Jin; GHIM, Sa Youl. Characterization of three antifungal calcite-forming bacteria, *Arthrobacter nicotianae* KNUC2100, *Bacillus thuringiensis* KNUC2103, and *Stenotrophomonas maltophilia* KNUC2106, derived from the Korean Islands, Dokdo and their application on mortar. *Journal of microbiology and biotechnology*, v. 23, n. 9, p. 1269-1278, 2013.

PATEL, Kartik; PATEL, Mitesh. Improving bioremediation process of petroleum wastewater using biosurfactants producing *Stenotrophomonas* sp. S1VKR-26 and assessment of phytotoxicity. *Bioresource Technology*, v. 315, p. 123861, 2020.

PATEL, Vilas; PATEL, Janki; MADAMWAR, Datta. Biodegradation of phenanthrene in bioaugmented microcosm by consortium ASP developed from coastal sediment of Alang-Sosiya ship breaking yard. *Marine pollution bulletin*, v. 74, n. 1, p. 199-207, 2013.

PATIL, Chandrashekar et al. Improved method for effective screening of ACC (1-aminocyclopropane-1-carboxylate) deaminase producing microorganisms. *Journal of microbiological methods*, v. 131, p. 102-104, 2016.

PATOWARY, Kaustuvmani et al. Development of an efficient bacterial consortium for the potential remediation of hydrocarbons from contaminated sites. *Frontiers in microbiology*, v. 7, p. 1092, 2016.

PERMALA, Ratish R. et al. Complete Genome Sequence of *Stenotrophomonas maltophilia* AB550, an Environmental Solar Radiation-and Multidrug-Resistant Strain Isolated in Western Australia. *Microbiology Resource Announcements*, v. 7, n. 5, p. e00914-18, 2018.

RADZUAN, Mohd Nazren; BANAT, Ibrahim M.; WINTERBURN, James. Production and characterization of rhamnolipid using palm oil agricultural refinery waste. *Bioresource Technology*, v. 225, p. 99-105, 2017.

RIBEIRO, Beatriz G.; GUERRA, Jenyffer MC; SARUBBO, Leonie A. Biosurfactants: Production and application prospects in the food industry. *Biotechnology progress*, v. 36, n. 5, p. e3030, 2020.

ROCHA E SILVA, Nathalia Maria P. et al. Natural surfactants and their applications for heavy oil removal in industry. *Separation & Purification Reviews*, v. 48, n. 4, p. 267-281, 2019.

RYAN, Robert P. et al. The versatility and adaptation of bacteria from the genus *Stenotrophomonas*. *Nature reviews microbiology*, v. 7, n. 7, p. 514-525, 2009.

SAHU, Kuleshwar Prasad et al. Rice blast lesions: an unexplored phyllosphere microhabitat for novel antagonistic bacterial species against *Magnaporthe oryzae*. *Microbial ecology*, v. 81, p. 731-745, 2021.

SANZ-GARCÍA, Fernando et al. Coming from the wild: multidrug resistant opportunistic pathogens presenting a primary, not human-linked, environmental habitat. *International Journal of Molecular Sciences*, v. 22, n. 15, p. 8080, 2021.

SANTOS, Danyelle Khadydja F. et al. Biosurfactants: multifunctional biomolecules of the 21st century. *International journal of molecular sciences*, v. 17, n. 3, p. 401, 2016.

SARUBBO, Leonie A. et al. Biosurfactants: Production, properties, applications, trends, and general perspectives. *Biochemical Engineering Journal*, p. 108377, 2022.

SHAH, Devendra H. et al. The occurrence of Salmonella, extended-spectrum β -lactamase producing Escherichia coli and carbapenem resistant non-fermenting Gram-negative bacteria in a backyard poultry flock environment. *Zoonoses and Public Health*, v. 67, n. 6, p. 742-753, 2020.

SHAHZAD, Asim; SADDIQUI, Samina; BANO, Asghari. The response of maize (*Zea mays* L.) plant assisted with bacterial consortium and fertilizer under oily sludge. *International journal of phytoremediation*, v. 18, n. 5, p. 521-526, 2016.

SHAHID, Muhammad Kashif et al. Current advances in treatment technologies for removal of emerging contaminants from water—A critical review. *Coordination Chemistry Reviews*, v. 442, p. 213993, 2021.

SHI, Juan et al. Rhamnolipid production from waste cooking oil using newly isolated halotolerant *Pseudomonas aeruginosa* M4. *Journal of Cleaner Production*, v. 278, p. 123879, 2021.

SHREYA, Desai et al. Amelioration effect of chromium-tolerant bacteria on growth, physiological properties and chromium mobilization in chickpea (*Cicer arietinum*) under chromium stress. *Archives of microbiology*, v. 202, p. 887-894, 2020.

SIDDIQUE, Tariq et al. Bacterial reduction of selenium in coal mine tailings pond sediment. *Journal of Environmental Quality*, v. 36, n. 3, p. 621-627, 2007.

SILVA, Rita De Cássia FS et al. Applications of biosurfactants in the petroleum industry and the remediation of oil spills. *International journal of molecular sciences*, v. 15, n. 7, p. 12523-12542, 2014.

SINGH, Paramjit; PARMAR, Nitika. Isolation and characterization of two novel polyhydroxybutyrate (PHB)-producing bacteria. *African Journal of Biotechnology*, v. 10, n. 24, p. 4907-4919, 2011.

SNELLINX, Zita et al. Microbial consortia that degrade 2, 4-DNT by interspecies metabolism: isolation and characterisation. *Biodegradation*, v. 14, p. 19-29, 2003.

SOHAIL, Rafeya; JAMIL, Nazia. Isolation of biosurfactant producing bacteria from Potwar oil fields: Effect of non-fossil fuel based carbon sources. *Green Processing and Synthesis*, v. 9, n. 1, p. 77-86, 2019.

SOMARAJA, P. K.; GAYATHRI, D.; RAMAIAH, N. Molecular characterization of 2-chlorobiphenyl degrading *Stenotrophomonas maltophilia* GS-103. *Bulletin of environmental contamination and toxicology*, v. 91, p. 148-153, 2013.

SONDH, Sidhartha et al. A strategic review on Municipal Solid Waste (living solid waste) management system focusing on policies, selection criteria and techniques for waste-to-value. *Journal of Cleaner Production*, p. 131908, 2022.

SUBRAMANIAN, Kumaran et al. Bioconversion of chitin waste through *Stenotrophomonas maltophilia* for production of chitin derivatives as a Seabass enrichment diet. *Scientific Reports*, v. 12, n. 1, p. 4792, 2022.

URSZULA, Guzik et al. Isolation and characterization of a novel strain of *Stenotrophomonas maltophilia* possessing various dioxygenases for monocyclic hydrocarbon degradation. *Brazilian Journal of Microbiology*, v. 40, p. 285-291, 2009.

WAGHMARE, Shailesh R. et al. Purification and characterization of novel organic solvent tolerant 98 kDa alkaline protease from isolated *Stenotrophomonas maltophilia* strain SK. *Protein Expression and Purification*, v. 107, p. 1-6, 2015.

WILSON, Mark; LINDOW, Steven E. Coexistence among epiphytic bacterial populations mediated through nutritional resource partitioning. *Applied and environmental microbiology*, v. 60, n. 12, p. 4468-4477, 1994.

XIONG, Weiliang et al. Characterization of an efficient estrogen-degrading bacterium *Stenotrophomonas maltophilia* SJTH1 in saline-, alkaline-, heavy metal-contained environments or solid soil and identification of four 17 β -estradiol-oxidizing dehydrogenases. *Journal of hazardous materials*, v. 385, p. 121616, 2020.

YOUENOU, Benjamin et al. Comparative genomics of environmental and clinical *Stenotrophomonas maltophilia* strains with different antibiotic resistance profiles. *Genome biology and evolution*, v. 7, n. 9, p. 2484-2505, 2015.

ZAFFAR, Habiba et al. Kinetics of endosulfan biodegradation by *Stenotrophomonas maltophilia* EN-1 isolated from pesticide-contaminated soil. *Soil and Sediment Contamination: An International Journal*, v. 27, n. 4, p. 267-279, 2018.

ZHANG, Weihong et al. Uniqueness and dependence of bacterial communities on microplastics: comparison with water, sediment, and soil. *Microbial ecology*, p. 1-11, 2021.

ZOU, Changjun et al. Characterization and optimization of biosurfactants produced by *Acinetobacter baylyi* ZJ2 isolated from crude oil-contaminated soil sample toward microbial enhanced oil recovery applications. *Biochemical engineering journal*, v. 90, p. 49-58, 2014.